2188

Discussion Papers

Deutsches Institut für Wirtschaftsforschung 2026

Patient-driven Information Flow and Practice Style Spillover

Temulun Borjigen

Opinions expressed in this paper are those of the author(s) and do not necessarily reflect views of the institute.

IMPRESSUM

© DIW Berlin, 2026

DIW Berlin
German Institute for Economic Research
Anton-Wilhelm-Amo-Str. 58
10117 Berlin

Tel. +49 (30) 897 89-0
Fax +49 (30) 897 89-200
http://www.diw.de

ISSN electronic edition 1619-4535

Papers can be downloaded free of charge from the DIW Berlin website:
http://www.diw.de/discussionpapers

Discussion Papers of DIW Berlin are indexed in RePEc and Econstor:
http://ideas.repec.org/s/diw/diwwpp.html
https://www.econstor.eu/handle/10419/10

Patient-driven Information Flow and Practice Style Spillover*

Temulun Borjigen†

October 2026

Abstract

How physicians form and update practice styles remains central to understanding persistent variation in care. Using quasi-random patient reallocation following primary care clinic closures in Denmark, I examine whether antibiotic prescribing practice styles are transmitted across clinics through patient mobility. I find that clinics receiving more patients with prior antibiotic use increase prescribing intensity among their existing patients. A twofold increase in incoming patients with prior antibiotic use raises prescribing intensity to existing patients by 2.3 percent. The response is stronger among newer clinics and is accompanied by increased use of diagnostic tests, consistent with physicians learning from the prescribing histories of incoming patients rather than responding only to patient volume. The findings highlight patient mobility as an important, informal channel of information transmission in healthcare markets and show that prescribing practices can diffuse across clinics through patient reallocation. These spillovers have implications for the persistence of practice variation and for the design and evaluation of policies aimed at improving antibiotic stewardship.

JEL Classification: I11, I12, I18, D83

Keywords: Practice style; Physician learning; Prescribing behavior; Patient mobility; Information diffusion; Antibiotic stewardship

*I thank Tomaso Duso, Hannes Ullrich, participants at the DIW Berlin IO Brown Bag Seminar, and members of the Health Care Markets and Health Policy Research Group at ZEW Mannheim for helpful comments and discussions. I also gratefully acknowledge financial support from the European Research Council under the European Union's Horizon 2020 research and innovation programme (grant agreement no. 802450).

†Department Firms and Markets, DIW Berlin; Department of Economics, University of Copenhagen; and Berlin School of Economics. tborjigen@diw.de

1 Introduction

Healthcare providers exhibit substantial and persistent variation in the quality and intensity of care they deliver. A large body of research documents wide differences in treatment choices, spending, and outcomes across physicians and healthcare markets, even for clinically similar patients (e.g., Chandra et al. 2011; Finkelstein et al. 2016; Fadlon and Van Parys 2020; Huang and Ullrich 2024). Such variation is difficult to reconcile with demand-side differences in patient preferences or medical need alone. It points instead to persistent supply-side heterogeneity and raise concerns about inefficiencies, unequal access to high-quality care, and uneven adoption of appropriate treatment practices.

A key supply-side driver for this variation is heterogeneity in physician beliefs about appropriate treatments (Phelps 1992; Cutler et al. 2019). Physicians may differ in how they interpret clinical evidence, assess treatment risks, or respond to diagnostic uncertainty. These differences stem in large part from uneven exposure to information regarding the availability and the efficacy of treatments, as well as best practices (Phelps 2000; Arrow et al. 2020). Understanding how physicians acquire information and update beliefs is therefore critical for evaluating policies aimed at reducing unwarranted variation and improving healthcare quality.

Physicians acquire information through multiple channels. Formal medical training and continuing education shape baseline knowledge, while clinical guidelines and decision-support tools provide standardized recommendations. Information may also diffuse informally through interaction with peers in team practices, professional networks, or referral relationships (Coleman et al. 1957; Chen 2021; Agha et al. 2022). Clinical databases and scientific publications also provide information generated by peer physicians, facilitating adoption of new technologies and prevailing standards of care (Epstein and Ketcham 2014; Agha and Molitor 2018; Arrow et al. 2020).

In this paper, I focus on a distinct channel of information transmission: patient mobility across primary care clinics. When patients move between providers, they bring with them realized treatment histories, including prior diagnoses, prescriptions, and responses to earlier care. These histories can inform receiving physicians in two ways. First, they provide evidence about previous treatment decisions and outcomes, which may reduce diagnostic or treatment uncertainty, particularly when symptoms persist or previous treatments appear ineffective. Second, they may reveal how peer physicians have treated similar cases, thereby transmitting information about prevailing practice styles in the local primary care environment. Unlike guidelines or formal communication, this information is generated through routine patient care and may affect decision-making even without direct interaction between providers.

I investigate whether this information channel affects physicians' decisions in the context of antibiotic prescribing in primary care, using administrative data covering the universe of patients and primary care clinics from 2014 to 2019 in Denmark. Antibiotic prescribing is a particularly informative setting because treatment decisions have clear quality and public health implications. Inappropriate prescribing, whether driven by patient demand, diagnostic uncertainty, or physician practice style, is a major contributor to antimicrobial resistance, a growing

global health concern (Laxminarayan et al. 2013). Broad-spectrum antibiotics are especially relevant because they are sometimes used when narrower alternatives would be clinically sufficient. Changes in antibiotic prescribing behavior therefore have direct welfare and policy relevance.

Furthermore, prescribing decisions are often made with limited diagnostic information (Manski 2018), requiring physicians to rely on experience and informal signals. In Denmark, primary care physicians act as gatekeepers of antibiotics, and account for 88% of the total antibiotics prescribed in 2024.[1] However, antibiotic prescribing exhibits large and persistent variation across physicians and clinics that is difficult to reconcile with differences in patients' medical need (Huang and Ullrich 2024). To promote rational and efficient use of antibiotics, the Danish Health and Medicines Authority issued a guideline[2] for physicians in 2013. Nevertheless, physicians often make prescribing decisions with incomplete diagnostic information, as tests for bacterial infection may be delayed or unavailable at the point of care (Ribers and Ullrich 2023). In these cases, physicians must rely on clinical judgment and prior experience to determine whether antibiotics are needed, and differences in diagnostic skill can generate systematic variation in treatment choices (Currie and MacLeod 2020). Hence, understanding how prescribing behavior forms and updates is essential for designing interventions that reduce unwarranted variation and improve the quality of antibiotic use.

Empirically isolating the effect of information transmitted through patient flows is challenging because patient movements are typically endogenous to physician behavior and clinic characteristics. To address this endogenous selection, I exploit quasi-random patient reallocation following primary care clinic closures in Denmark. Clinic closures reassign affected patients to nearby clinics, generating plausibly exogenous variation in both the number and composition of incoming patients across receiving clinics. Using Danish administrative data covering the universe of patients and primary care clinics from 2014 to 2019, I identify clinic closures and subsequent patient switches, and construct a shift-share style instrument that captures receiving clinics' exposure to the prescribing styles of closed clinics through patient arrivals. Specifically, the instrument weights the antibiotic prescribing style of the closed clinics, defined as the average prescribing intensity to their existing patients, by each clinic's share of the displaced patients it receives. This instrument isolates variation in the antibiotic prescribing volume of newly arrived patients that is driven by their origin clinics' styles rather than by the receiving clinic's own practices or patient selection.

To abstract from changes in prescribing that reflect the medical needs of incoming patients, I restrict the outcome to the existing patient base of the receiving clinic, including only patients who are listed before the closure-induced inflow and do not switch clinics. I measure the receiving clinic's information exposure through arriving patients' antibiotic prescribing history, defined as at least one antibiotic prescription preceding their origin clinic's closure. Instrumenting this

[1] Summary Report 2024, The Danish Integrated Antimicrobial Resistance Monitoring and Research Programme (DANMAP), https://www.danmap.org/-/media/institutter/foedevareinstituttet/danmap-site/report-2024/summary_danmap_2024_full-report_low-vesion-2.pdf

[2] Guidelines on prescribing antibiotics for physicians and others in Denmark, the Danish Health and Medicines Authority, https://www.sst.dk/media/4b5het54/guideline-on-prescribing-antibiotics.pdf. Last accessed 13/01/2026.

inflow of information with the shift-share exposure to closed clinics' prescribing styles, I find that receiving double the number of patients with an antibiotic prescribing history increases the overall antibiotic prescribing intensity for existing patients at the receiving clinic by 2.3 percent. The effect is larger among newer clinics, consistent with models of Bayesian learning in which agents with less precise prior information respond more strongly to new signals.[3] However, this increase does not come at the cost of care quality and access, as I find positive effects on both diagnostic testing and prescriptions issued before test results are available, but no evidence of an increase in prescriptions that require subsequent correction. These patterns suggest that the response is not driven solely by mechanically more aggressive or lower-quality prescribing.

By documenting how prescribing behavior responds to information embedded in patient histories, this paper contributes to the literature on physician learning and information acquisition. A growing body of work shows that physicians update beliefs based on accumulated experience and imperfect signals, particularly in the context of new or generic drug diffusion. These signals include responses from their own patients as well as information recorded in administrative or clinical databases on treatment outcomes for other patients (Coscelli and Shum 2004; Epstein and Ketcham 2014; Arrow et al. 2020; Zhu 2023). Related learning processes also occur on the demand side, where patients update beliefs about drug effectiveness through their own experience, social networks, and word of mouth (Crawford and Shum 2005; Ching 2010). A separate literature on volume-outcome relationships shows that physicians and hospitals performing more procedures often achieve better outcomes, consistent with learning from repeated practice (Gaynor et al. 2005; Avdic et al. 2019; Domenella 2025). This literature primarily emphasizes learning from within-provider experience. Evidence is lacking on whether and how providers learn through exposure to patients who have previously been treated by other physicians.

This paper also relates to the literature that has documented important variation in healthcare provision across providers. Prior studies document large and systematic differences in treatment intensity and quality at the provider level across geographic areas (Chandra and Staiger 2007; Finkelstein et al. 2016). Other work shows substantial within-region variations in care delivery, driven in part by provider-specific practice styles (Fadlon and Van Parys 2020; Huang and Ullrich 2023), which can be as large as, or larger than, variation across regions (Epstein and Nicholson 2009). While these studies establish the importance of provider-specific factors, fewer papers examine how such practice styles are formed and transmitted. Existing evidence points to environmental and regional factors, including hospital capacity, productivity, and local policies (Chandra and Staiger 2007; Epstein and Nicholson 2009; Molitor 2018). Other studies have shown that financial incentives and market competition can shift treatment decisions, without explicitly translating the observed effect into practice styles (Currie et al. 2014; Bennett

[3] For example, Rockoff et al. (2012) finds that school principals with less precise prior beliefs change their evaluation of teacher productivity more after receiving an external performance measure of the teachers. Bhuller and Sigstad (2025) shows that after learning reversals in appeal decisions, less experienced judges and judges who have not processed a similar case before react more by adjusting their sentencing for future cases. In the healthcare context, Coscelli and Shum (2004) and Crawford and Shum (2005) show that prior beliefs about drug effectiveness are refined over time once physicians and patients start to observe the treatment outcomes. Although they did not explicitly parameterize updating by physician experience, both models suggest that responses to new signals depend on the priors accumulated over time.

et al. 2015). In this paper, I show that incoming patients can transmit information about prior prescribing practices across clinics, thereby shaping treatment decisions at the receiving clinic. By transmitting information between providers, moving patients may be a factor in reducing persistent variation in providers' practice styles without the need for explicit standardization.

Finally, this paper relates to the broader literature on peer effects and knowledge spillovers. Previous work shows that physician behavior responds to interactions with peers, referral networks, and local professional environments (Coleman et al. 1957; Chen 2021; Agha et al. 2022). Similar spillover mechanisms have been documented in other social settings. For example, Sorensen (2006) finds that coworkers learn about their healthcare plans and converge towards the same choices. Another example from Bailey et al. (2022) in the consumer market shows that people are more likely to purchase phones of the same brand following a recent purchase by their friend. This paper extends the peer effects literature by demonstrating that knowledge spillovers need not operate through direct communication or social networks. Instead, patients can transmit information across providers through their treatment histories, allowing physicians to infer peer practices indirectly. This mechanism complements existing evidence on learning through professional interaction and highlights patient mobility as an additional channel through which provider behavior can evolve.

2 Institutional Background and Data

2.1 Institutional Background

Danish public health insurance has universal coverage of visits and services from primary care providers, who are usually general practitioners (also known as family doctors). General practitioners can take care of minor health issues and health maintenance. Additionally, the general practitioner also acts as a gatekeeper for other healthcare services such as prescriptions for drugs, referrals to other specialists, and hospital inpatient treatments. There are about 3,500 general practitioners in Denmark, each treating about 1,600 patients (Simonsen et al. 2021). In 2019, approximately 1,700 clinics were run by general practitioners, half of which were single-physician clinics.[4] Each primary care provider has to acquire a practice authorization number to claim reimbursement from public health insurance. The state has control over the number of practice authorizations based on factors such as the population density in a given area.

Primary care providers are self-employed and are responsible for the revenue of their clinics. A clinic's income is composed of two parts: a fixed fee per listed patient, i.e. capitation, which accounts for about one-third of their income, and fee-for-service payments that make up the rest. Physicians do not have agency over the fees they charge. Instead, the Danish Medical Association and the government negotiate the charges every third year (with biannual price adjustments). Notably, physicians do not receive additional fees directly related to prescriptions.

[4] Praktiserende Lægers Organisation (PLO), *PLO-læge- og praksispopulationen 1977–2022: Nøgletal fra medlemsregisteret*, Table 1.6, p. 40, https://laeger.dk/media/3kkkkp1l/laege-_og_praksispopulationen_1977-2022.pdf (accessed July 26, 2026). The statistics cover physicians who are PLO members and attached to a provider number.

Every resident in Denmark is required to join the patient list of a primary care clinic, typically located within a 15 km radius of their residence[5], which will serve as their designated contact in case of medical needs. Physicians cannot selectively reject individual patients as long as they have not reached a capacity limit of 1,600.[6] Physicians can stop accepting new patients when they have 1,600 patients on their list. Patients can change their initially listed practice for a fee (DKK 150, about USD 20). Only the physician's name and the clinic address are publicly observed, so that selective switching between primary care clinics is difficult (Huang and Ullrich 2023).

In the case of a closure, a new general practitioner can acquire the entire patient list, often with a takeover of the physical practice. If no such takeover occurs, patients of the closing clinics are randomly assigned to nearby practices that are accepting new patients. Closures do not need to be communicated to the patients in advance, but patients are notified of clinics that are open for new patients within the 15 km limit, and the local government is obliged to ensure that at least two practices are available to be chosen. When patients switch to a new clinic, their health records are transferred automatically, unless they explicitly deny the transfer within the first 14 days after the switch. This feature is crucial for the identification of the information transmission, as it guarantees that the receiving clinics have access to both documented and self-communicated treatment histories from switching patients.

2.2 Data and Sample Construction

Clinics. I use authorization numbers from the Danish Clinic Registry (Yderregisteret), matched to claims data from the Danish National Outpatient Claims Registry (Sygesikringsregistret), to identify primary care clinics from 2014 to 2019. The Clinic Registry reports, for each clinic, opening and closing dates, postcode, and the unique personal identification numbers (personnummer, PNR[7]) of physicians and other healthcare workers affiliated with the clinic. The Claims Registry contains clinic authorization numbers, the date a claim is filed, patient identifiers (PNR), and the types of medical services provided. Medical services are recorded using six-digit codes, which allow me to identify services that are exclusively provided by the primary care sector.

The outcome variables and prescribing histories are constructed using prescription data from the Danish National Prescription Registry (Lægemiddeldatabasen, LMDB). This registry records every prescription redeemed at Danish pharmacies, including the drug name, brand, quantity, dispensing date, clinic authorization number, and patient PNR. I then link the Clinic Registry, claims data, and prescription data using the clinic authorization numbers to construct a clinic-level panel of claims and prescribing activity. I further use this panel to infer missing information for some clinics. For example, for clinics without a recorded closing date in the Clinic Registry, I assume that they remain active until there is a sharp and sustained decline in prescription volumes, typically accompanied by a similar decline in claims. Through match-

[5] 5 km in densely populated areas such as central Copenhagen.

[6] In some cases, this capacity limit may not be binding so that some physicians have over 1,600 patients on their lists.

[7] The PNR is the unique pseudonymized identifying number assigned to all residents of Denmark.

ing clinic activities, I obtain a preliminary sample of 2,207 primary care clinics with complete information on clinic activities, characteristics and patient compositions from 2014 to 2019.

Patient-provider assignment. I define each patient's choice of primary care clinic and identify switching behavior following the algorithm proposed by Kjaersgaard et al. (2016). The algorithm uses general practice service records from the Claims Registry to assign patients to clinics over time based on observed patient–provider encounters. Specifically, patients are linked to the clinic providing their general practice services, and continuous patient–clinic affiliation intervals are constructed using the timing and provider identifiers of these services. Very short affiliation spells (less than 31 days) are removed to focus on sustained relationships, and the resulting intervals are aggregated to 3 calendar months to create a consistent quarterly panel, where each patient is assigned to exactly one primary care clinic.[8] A patient is defined as switching clinics when the assigned clinic changes from one quarter to the next. This procedure closely replicates the official Patient List Database[9] and provides a reliable measure of patient choice and switching over time.

Estimation sample. Since patient movements can only be identified through two consecutive quarters, for the estimation sample, I drop the first observation for every clinic. Then I exclude transition periods where the destination clinics potentially take over the closed clinics, as clinics that replace an existing clinic do not have an existing pool of non-switching patients to study their outcomes.[10] I assume a new clinic is a takeover when it is at the same location as a closed clinic, and its entry date is in the same or next quarter in which the closure happens. For these takeovers, I drop the periods where they receive more patients from closed clinics than non-switchers. Similarly for both existing and other new clinics that are not takeovers, I remove the periods where they do not have a sufficiently large number of non-switching patients to identify any changes to their prescribing patterns.[11] Finally, to reduce noise arising from sparse observations and to avoid mechanically large ratios driven by small denominators, I exclude observations with very small base values. When constructing share-based measures such as the prescribing intensity and the share of patients with prior prescriptions among newly received patients, I drop clinic-quarter period observations with a low number of consultations and new patients.[12]

[8] I drop weeks with services from multiple clinics to avoid ambiguous assignments, in line with Kjaersgaard et al. (2016).

[9] The quarterly patient list constructed using this procedure shows 93.4% agreement with the official data. Kjaersgaard et al. (2016) reports higher overall match of 98.6% for monthly patient lists.

[10] As a robustness check, I also estimate the main regression on two subsamples. First, I exclude all observations of clinics that are takeovers and for the other I exclude all newly entered clinics between 2014 and 2019. The results are in Appendix Table E.11.

[11] Specifically, I drop clinic-quarter observations in the bottom 5% in terms of the number of non-switching patients.

[12] For the number of consultations from the non-switchers, I drop observations with fewer than 50 consultations per quarter (bottom 5%). I also drop quarters where the clinic receives fewer than 5 new patients to reduce noise as well as to ensure anonymity.




Figure 2.1: Number of primary clinics and average clinic size

Notes: This graph illustrates the development in the number of primary care clinics and clinic size in Denmark, during 2014-2019. Clinic size is defined as the number of physicians actively practicing in a given quarter, excluding other supportive roles such as assistants, administrative staff, and interns. Primary care clinics are considered active if they file claims and issue a reasonable number of prescriptions. Clinics that did not file any claims in a given quarter are considered inactive and are dropped.

2.3 Descriptive Statistics

My final sample contains 5,011,537 patients assigned to 1,838 active primary care clinics from 2014 to 2019. The majority of the clinics in the sample (87%) were established before the first quarter of 2014, the earliest sample date. Among all clinics opened before the first quarter of 2014, 1,152 remained active until the final quarter of 2019. [13] During the sample period, 374 clinics closed permanently and 248 new clinics opened, which is consistent with the nationwide decreasing trend in the number of primary care clinics shown in Figure 2.1. However, the average number of physicians in a clinic was increasing in the sample period.

Table 2.1 summarizes descriptive statistics of the clinic-quarter sample. A primary care clinic has on average 2.2 physicians and around 3,100 listed patients. The average number of 1,600 patients per physician corresponds to what the previous literature (e.g. Haastrup et al. 2025) has documented, supporting the quality of the matching algorithm. During 2014 to 2019, an average clinic filed around 8,800 claims per quarter, 28% of which were consultations. I distinguish consultations from other types of medical services because they serve as a proxy for the frequency of patient visits, and also because the majority of antibiotic prescriptions in primary care are given during consultations. Other types of treatments that primary care clinics carry out and file claims for include referrals to specialists and follow-ups of the previous treatment. Variation in the number of claims and consultations across clinics is mostly due to variation in clinic size as, by design, bigger clinics with more physicians have a larger capacity to admit patients. Hence, when analyzing primary care clinics' medical service provision, I net out the effect of clinic size or the number of patients by averaging the total number of prescriptions over the number of consultations at a given time and clinic.

[13] 50% of these 1,152 clinics had complete records for every quarter. Missing clinic activities are primarily due to temporary closures of the clinics.

Table 2.1: Clinic descriptive statistics

	mean	s.d.
Number of clinics	1,399.87	96.23
Total number of listed patients	3,078.23	1,848.15
Number of visited patients	2,308.67	1,492.47
Number of antibiotic prescriptions	296.73	200.87
Defined daily dose (DDD)	2,995.73	2,026.52
Number of claims	8,782.97	5,758.08
Number of consultations	2,919.05	1,857.91
Number of prescriptions per consultation	0.10	0.04
Number of claims per physician	4,152.76	1,404.94
Clinic size	2.18	1.34
Share of female physicians	0.49	0.40
Physician age	53.66	7.84
Physicians with immigration background	0.08	0.24
Years since entry	21.32	14.18
Non-switching patients		
Number of non-switching patients	2,118.54	1,663.17
Share of non-switching patients	0.65	0.26
Number of antibiotic prescriptions	173.28	153.90
Number of consultations	1,856.51	1,578.89
Number of claims	5,480.88	4,719.56
Number of prescriptions per consultation	0.10	0.04
Other patients		
Number of new patients	52.02	66.91
Share of new patients with prescriptions	0.30	0.13
Number of prescriptions per consultation	0.12	0.05
Observations	32,197	

Note: This table summarizes the mean and the standard deviation of the listed variables from an unbalanced quarterly panel consisting of 1,838 active primary care clinics from 2014 to 2019. Clinic size refers to the number of physicians in a clinic, excluding other supportive roles such as assistants, administrative staff, and interns. Physicians with an immigration background include physicians who are first- or second-generation immigrants.

Table 2.2: Share of patient types


Non-switchers	2,357,826	47%
Switchers	2,653,711	53%
Switch due to clinic closure	919,424	18%
Switch due to move	732,412	15%
Switch due to other reasons	1,001,875	20%

The bottom two panels of Table 2.1 report descriptive statistics at the clinic level, distinguishing between non-switching and new patients. Clinics serve a large and stable base of non-switching patients, with an average of around 2,100 such patients per clinic and a mean share of 65% of the patient population. This stability is also reflected in high levels of patient activity, where non-switching patients account for an average of 1,856 consultations and 173 antibiotic prescriptions per clinic. Despite substantial variation across clinics, as indicated by the large standard deviations, prescribing intensity for non-switching patients is relatively stable, with an average of 0.10 prescriptions per consultation.


In contrast, clinics receive a much smaller inflow of new patients. On average, clinics see 52 new patients per quarter, who constitute a considerably smaller share of the patient population. Among these incoming patients, 30% have a prior history of antibiotic prescriptions. The substantial variation in both the number of new patients and their prior prescribing histories generates heterogeneity in the information arriving at receiving clinics. Once they are admitted, new patients are more likely to receive a prescription conditional on a consultation, with a mean of 0.12 prescriptions per consultation, compared to non-switching patients.

Table 2.2 summarizes the switching decisions among patients in the sample. Switchers are patients who have changed their primary clinics at least once based on the matching algorithm described in Section 2.2. Non-switchers are patients who remain with the same clinic during the sample period. More than half (53%) of the patients moved to a different clinic at least once in 6 years between 2014 and 2019, and one-third of them switched because their previous clinics had closed. Another third of the switchers changed clinics because they moved to different home addresses. Appendix Table B.1 compares the characteristics of switching and non-switching patients. While switchers, on average, receive fewer antibiotic prescriptions and have fewer consultations per person, the two groups are otherwise broadly comparable in terms of healthcare utilization and demographic characteristics. There are higher shares of female patients and patients with an immigration background among switchers, and patients who switch are also younger, with a mean age of 38 at the time of their switches.

I further present two stylized facts that characterize switching patients and their receiving clinics in Figure 2.2. First, the share of new patients with prior antibiotic prescribing is proportional to the number of new patients at the receiving clinic. As shown in Panel (a), doubling the number of new patients increases the likelihood of receiving a patient who comes with a prescribing history by 2 percentage points. In other words, the more new patients that come to a



(a) Relationship between number of new patients and share of prescribed new patients



(b) Relationship between lagged prescribing intensity and number of new patients

Figure 2.2: Arrival of new patients and information at the receiving clinics

Notes: These figures present a binned scatter plot of the residualized outcome and residualized (log-)number of new patients. Both variables in each figure are residualized with respect to clinic and year-quarter fixed effects prior to binning. The figure then bins the clinic-level observations into 20 equal-sized bins based on residualized (log-)number of new patients. Each dot represents the mean of the outcome within a bin of the explanatory variable. The solid line shows the fitted linear relationship based on the underlying data.

clinic, the more likely the clinic is exposed to information specific to the prescribing practices of other clinics. Second, I show that the arrival of new patients is not correlated with the receiving clinic's past prescribing intensity in Panel (b). There is a small and statistically insignificant negative correlation between prescribing intensity in the period before patient arrival and the number of new patients. These figures, together with the decomposition of reasons for switching presented in Table B.1, lend confidence to the assumption that switching is likely triggered by factors exogenous to a clinic's performance rather than selective sorting based on prescribing behaviors.

Appendix Figure B.1a plots the distribution of the changes in distances to the clinics after switching to a new clinic. The majority of patients move to clinics closer than their original

clinics, and patients are more likely to switch within the vicinity of their home address, as shown by the concentration around a 1 km difference in distance. Furthermore, Appendix Figure B.1b suggests that patients' choice of clinic does not appear to be dependent on prescribing patterns, and if anything, more patients switch to clinics with lower average prescribing intensities than their original clinics. This additional evidence again supports that patients do not actively sort on the prescribing behaviors of the receiving clinics.

3 Empirical strategy

3.1 Measures for exposure to information

To estimate the effect of information from new patients on antibiotic prescribing at the receiving clinic, I define the main information measure, *volume*, using the incoming patients and their prescribing histories. The measure counts the number of new patients who received *any* prior antibiotic prescriptions at their origin clinics.[14] Information about past treatment may reach the receiving physician through electronic health records, which are transferred when patients switch clinics, or directly through communication with the patient. The volume measure therefore captures the potential amount of prescribing information available to the receiving clinic and represents the extensive margin of information exposure: clinics that receive more new patients with prior prescriptions observe a larger number of prescribing signals. Formally, for each receiving clinic in each quarter, I measure the size of the information inflow by the number of new patients who received *any* number of antibiotic prescriptions at their origin clinics before switching:

$$volume_{it} = \sum_{j=1}^{J} (prescribed_{jt} \times switch_{ijt})$$

Here, $prescribed_{jt} \in \{0, 1\}$ is an indicator for patient j receiving an antibiotic prescription from their previous clinics before switching at t, and $switch_{ijt} \in \{0, 1\}$ takes the value 1 if patient j switched to clinic i at time t and 0 otherwise.[15]

For the main specifications, I measure the information available to clinics by the number of incoming patients with a prior prescription. To complement this volume-based measure, I also examine the intensity of information exposure, defined as the share of all new patients received by a clinic in a given quarter who had been prescribed the treatment by their previous physicians. This measure captures the concentration of prescribing histories in the incoming patient pool while abstracting from factors that determine the scale of patient inflows, such as the size of the closing clinic and the receiving clinic's available capacity. Two clinics may receive the same number of previously prescribed patients but differ substantially in this share because their total

[14] As older prescriptions may be less salient for current treatment decisions, in Section 4.4 I discuss alternative measures that restrict prescribing histories to prescriptions observed within more recent quarters preceding the patient's switch.

[15] Note that the identity of the origin clinic is irrelevant for information inflow, which is assumed to be fully captured by the prescription history indicator.

inflows differ, exposing them to different informational environments. Conversely, clinics with the same share may receive very different numbers of prescribing histories. Conditional on the total number of new patients, the share therefore isolates the composition of the inflow, whereas the main measure captures its volume. I also discuss and present the results using alternative measures for information in Section 4.4 and Appendix Tables E.1-E.7.

3.2 Outcomes for antibiotic prescribing

The main outcome is antibiotic prescribing intensity, defined as the number of antibiotic prescriptions per consultation at the clinic level. Scaling prescriptions by consultations captures a clinic's prescribing behavior conditional on patient contact and separates changes in prescribing practice from mechanical changes in the volume of care. To further separate changes in physicians' prescribing behavior from the direct antibiotic demand of incoming patients, I construct the outcome using only patients who remain with the same clinic throughout the observational period (*non-switchers*). Focusing on these existing patients ensures that changes in prescribing intensity reflect changes in how clinics treat their incumbent patient population rather than the prescribing histories or treatment needs of newly arriving patients. This restriction also gives the estimated coefficient β a direct interpretation as the information spillover from incoming patients to prescribing decisions for existing patients.

In addition to overall prescribing intensity, I examine whether information spillovers affect the quality of antibiotic prescribing and treatment procedures. I consider two complementary dimensions. First, I distinguish between broad- and narrow-spectrum antibiotics, which provide information on the composition of antibiotic treatment and are commonly used to assess prescribing quality. Second, I analyze three treatment process indicators, including the intensity of diagnostic testing, the frequency of prescriptions issued without prior testing, and the incidence of follow-up prescriptions. These measures capture whether changes in prescribing are accompanied by changes in diagnostic effort and subsequent treatment decisions. As with the main prescribing outcome, all measures are constructed using the clinic's existing patient population (*non-switchers*), so that they reflect changes in physicians' treatment of incumbent patients rather than the direct treatment needs of incoming patients. Together, these outcomes allow me to assess whether the transmission of prescribing practices through patient inflows extends beyond the overall propensity to prescribe and affects how antibiotics are selected and how treatment decisions are made.

Following the classification from the Annual Epidemiological Report on Antimicrobial consumption in the EU/EEA [16], I classify antibiotic prescriptions into broad- and narrow-spectrum antibiotics. Appendix Table A.1 provides the full list of antibiotics and corresponding ATC codes used to construct these measures. Narrow-spectrum antibiotics target a more limited range of bacteria and are generally preferred when the likely cause of an infection can be identified. Broad-spectrum antibiotics act against a wider range of bacteria and are more commonly used

[16] European Centre for Disease Prevention and Control, https://www.ecdc.europa.eu/sites/default/files/documents/antimicrobial-consumption-ESAC-Net-annual-epidemiological-report-2023_0.pdf. Last accessed July 26, 2026.

when there is greater uncertainty about the appropriate treatment. Their use is associated with a higher risk of contributing to antimicrobial resistance, so clinical guidelines generally recommend narrow-spectrum antibiotics when appropriate. Therefore, the composition of prescribing provides a useful indicator of prescribing quality, with a greater reliance on narrow-spectrum antibiotics typically reflecting more targeted antibiotic use.

The interpretation of changes in antibiotic composition also depends on how physicians arrive at the treatment decision. One channel through which prescribing histories may affect current treatment is by providing information about how similar symptoms were treated in the past. For example, a newly arrived patient may report that a similar episode was previously treated successfully with antibiotics, or the physician may observe this treatment in the patient's medical record. Even when the prior prescription occurred several quarters earlier, such information may affect the physician's beliefs about the appropriate treatment of similar cases and reduce the perceived need for additional diagnostic testing when treating other patients with comparable symptoms.

I therefore complement the antibiotic-type outcomes with three measures of the treatment process: diagnostic testing intensity, prescriptions issued without prior testing, and follow-up prescriptions. Diagnostic tests include rapid tests, microscopic examinations, and bacterial cultures related to potential bacterial infections. I classify an antibiotic prescription as issued without prior testing when it cannot be linked to any such test within seven days before or after the prescription date.[17] I define a follow-up prescription as a prescription for an antibiotic from a different ATC3 subgroup within seven days of the initial prescription. Such changes in treatment may indicate that the initial antibiotic was a poorer match for the underlying infection. Together, these outcomes capture diagnostic effort and the quality of the initial treatment choice, providing additional evidence on whether information exposure changes how physicians make prescribing decisions.

3.3 Linear model of antibiotic prescribing

I first estimate the following model:

$$y_{it} = \beta volume_{it} + \delta X_{it} + FE_i + FE_t + \epsilon_{it}$$

The parameter of interest is β, which measures the effect of exposure to information on antibiotic prescribing outcome y_{it} measured for non-switchers. X_{it} includes a set of time-varying control variables. All regressions include year-quarter fixed effects to control for time-varying factors such as seasonality in bacterial infections, as well as clinic fixed effects to control for time-invariant unobservable characteristics of individual clinics.

Ideally, patients would choose clinics at random, generating exogenous variation in clinics' exposure to patient-specific information. In practice, patient choices may depend on perceived clinic characteristics, including prescribing tendencies. For example, patients with prior prescrib-

[17] I allow for a seven-day window because physicians may submit claims for diagnostic tests with a delay. Longer delays may lead some tested prescriptions to be classified as prescriptions without testing.

ing histories may select high-prescribing clinics, mechanically generating a positive correlation between informational exposure and subsequent prescribing. Selection on unobserved care quality correlated with prescribing may generate similar bias. Clinics that accept new patients may also differ systematically from other clinics: less congested clinics may diagnose more thoroughly and prescribe less, or may prescribe more to retain patients. These forms of selection can therefore bias OLS estimates of β in either direction.

Identification is preserved if patients do not observe clinic-specific prescribing patterns when choosing clinics, such that the prescribing history of arriving patients, $prescribed_{jt}$, is orthogonal to unobserved clinic shocks ϵ_{it}. Under this assumption, exposure to patients' prescribing histories provides valid variation for identifying β. In the next section, I exploit variation in the informational content of patient inflows following exogenous displacement to address remaining concerns about endogenous patient selection.

3.4 Constructing the instrument from clinic closures

To identify the causal effect of exposure to information from new patients on antibiotic treatment decisions for existing patients, I use an instrumental variable strategy that addresses endogenous patient selection based on clinics' prescribing styles. The strategy exploits primary care clinic closures, after which displaced patients are administratively reassigned to new clinics. Because closures are predominantly driven by physician retirement, they are unlikely to reflect unobserved local conditions captured by ϵ_{it}. The resulting reallocation of patients therefore generates an exogenous demand shock to remaining clinics and provides variation in their exposure to incoming patients' prescribing histories.

Patients arriving from closed clinics are endowed with the treatment styles of their original clinics that impact their realized prescription histories. For example, patients from high-prescribing style clinics are more likely to have received antibiotic prescriptions than patients from low-prescribing clinics. Exploiting this correlation between practice style and realized prescribing, I instrument for the receiving clinic's information exposure using a share-weighted mean of the practice styles of closing clinics. The share s_{ikt} is the fraction of closure-displaced patients received by clinic i in quarter t who originate from closing clinic k, among all closure-displaced patients received by clinic i in that quarter:

$$s_{ikt} = \frac{\sum_{j=1}^{J} switch_{ijkt}}{\sum_{k=1}^{k \in G_t} \sum_{j=1}^{J} switch_{ijkt}},$$

where G_t denotes the set of clinics that close in $t-1$. I include displaced patients who arrive at a receiving clinic within four quarters after the closure of their original clinic. Throughout the main analysis, I define practice style as the mean of antibiotic prescribing intensity, the ratio of prescriptions to consultations, among non-switchers over all periods in which clinic k is observed:

$$w_k = \frac{1}{|T_k|} \sum_{t \in T_k} \frac{Prescriptions_{kt}^{NS}}{Consultations_{kt}^{NS}},$$

where T_k denotes the set of periods in which clinic k is observed, $|T_k|$ is the number of such periods, and the superscript NS indicates that prescriptions and consultations are measured among non-switchers. Since non-switchers account for a large and stable proportion of a clinic's patient pool, as shown in the summary statistics Table 2.1, treatment decisions for the non-switching patient population should reflect primary care physicians' persistent, underlying practice styles. [18]

Conditional on a set of control variables accounting for clinics' patient mixes, patients' prescription histories reflect variations in the practice styles patients experience during their time with these closing clinics. At the same time, clinic closures generate plausibly exogenous shifts in patients' switching decisions. The instrument thus isolates the component of information inflow composition driven by external structural shocks due to closures from inflows driven by endogenous factors such as selective patient re-sorting. Resembling a shift-share design, I define the instrument as

$$z_{it} = \sum_{k=1}^{k \in G_t} s_{ikt} w_k,$$

where G_t denotes the set of clinics that close in $t-1$, so that patients from closed clinic k arrived at clinic i in period t, w_k is the practice style of closed clinic k, and s_{ikt} represents the share of patients j from clinic k arriving at clinic i.

Therefore, z_{it} captures the share-weighted exposure of the receiving clinics to the closed clinics' practice style, driven by the redistributed patients following a clinic closure. In Figure 3.1, I show that this share-weighted practice style instrument is strongly and positively correlated with the number of previously prescribed patients arriving at the receiving clinics. Patients coming from high-prescribing style clinics are thus more likely to carry a non-zero prescription history, and the receiving clinics, on the other hand, are more exposed to non-zero prescribing signals when they receive a higher share of patients from high-prescribing clinics.

The instrument can be interpreted as valid if the correlation between origin clinic closures and prescribing outcomes at the receiving clinics operates only through the exogenous reallocation of patients from clinics with differing prescribing styles, and not through other contemporaneous shocks to the receiving clinics' prescribing decisions. One threat to identification is that clinic closures may not be exogenous to local demand conditions or to features of the primary care market that also shape prescribing behavior. For example, closures may be more likely in areas with population decline, changes in age structure, or temporary increases in infection rates. In such cases, both the inflow of patients from closing clinics and the prescribing behavior of receiving clinics may respond to the same local trends. This would bias both OLS and IV estimates. To mitigate this concern, I include clinic and time fixed effects and control for observable patient and physician characteristics, which absorb part of the common variation in

[18] Apart from this static measure for prescribing style, I also conduct the same set of analyses by constructing a time-varying measure to allow for the potential evolution of prescribing styles over time. Specifically, I compute a moving average of prescribing intensity to the non-switchers within 4 quarters, and construct the instrument using the same set of incoming patient shares for the receiving clinics. The estimates in Appendix Table E.10 show that the effects are comparable in size and statistical significance to the time-invariant practice style definition.




Figure 3.1: First stage regression of endogenous information measure on IV

Notes: The figure presents a binned scatter plot of the residualized information measure and residualized IV. Both variables in the figure are residualized with respect to the set of controls included in the full model, and clinic and year-quarter fixed effects prior to binning. The figure then bins the clinic-level observations into 20 equal-sized bins based on residualized IV. Each dot represents the mean of the outcome within a bin of the explanatory variable.

local demand.

A related concern is that closures may reflect unobserved supply-side factors. For instance, closures may be more common among clinics that face staffing shortages, financial pressure, or stronger competition from nearby providers. Closures may also be more likely among clinics with unusually low prescribing tendencies if such clinics are less able to retain patients in competitive markets. In that case, patient reallocation after closure would reflect not only a movement of patients, but also a change in the prescribing styles present in the local market. The estimated effect of incoming patients could then partly capture how receiving clinics respond to a change in local market composition, rather than only the effect of learning from the prescribing histories of new patients. While the data do not record the reason for closure, physicians in closing clinics are on average 62 years old in my sample, which suggests that retirement is a likely explanation in many cases.

A further caveat is that closure shocks may coincide with broader structural or technological changes that also affect prescribing behavior across clinics. Closures may occur alongside regional primary care reforms or local efforts to improve antibiotic stewardship. If such changes influence both clinic survival and prescribing behavior, the instrument may partly capture these broader shocks rather than only the effect of patient inflows. In the Danish setting, however, this concern is less likely. Denmark introduced its antimicrobial resistance monitoring program in 1995, and the most recent national guideline on antibiotic prescribing was published in October 2013, several months before the start of the analysis period in 2014. This timing makes it less likely that closures after 2014 were driven by major contemporaneous policy changes focused on antibiotic prescribing.

To assess the potential relevance of these concerns, I perform balance tests by regressing the lagged antibiotic prescribing intensity, as well as other clinic activities, on the instrument. The

Table 4.1: Estimates of information effect on prescribing intensity

	(1) OLS	(2) OLS
Number of prescribed patients	0.0006	0.0008
	(0.0019)	(0.0018)
Clinic covariates	No	Yes
Patient covariates	No	Yes
Clinic FE	Yes	Yes
Year-quarter FE	Yes	Yes
Observations	32,197	32,197
Outcome mean	-2.36	-2.36
R-squared	0.77	0.77

Notes: Number of prescribed patients and the outcome, antibiotic prescriptions per consultation, are in log-scale. Clinic covariates include clinic size, average physician age, share of female physicians, and share of physicians with an immigration background. Patient covariates include average patient age, share of children (age <18), share of female patients, and share of patients with an immigration background. Standard errors in parentheses are clustered at the receiving clinic level.

results in Appendix Table C.1 show the absence of significant pre-trends. I also correlate clinic-level, predetermined variables with the instrument. Appendix Table C.2 shows that the receiving clinics are balanced across these variables. Finally, I test whether closures are correlated with clinic activities and characteristics that may affect prescribing styles. Appendix Figure D.2 shows that, although physician age and clinic location predict closure events, none of the other estimates are statistically significant. These results suggest that closure shocks are unlikely to be correlated with antibiotic prescribing behaviors.

4 Results

4.1 Effect on prescribing intensity

Table 4.1 reports the OLS relationship between the prescribing histories of incoming patients and prescribing among the receiving clinic's existing patients. Across specifications, the estimated coefficients are small and statistically indistinguishable from zero, providing little evidence of a systematic association in the raw observational relationship. Under voluntary patient choice, incoming patients' prescribing histories are potentially correlated with unobserved characteristics of the receiving clinic, as both patient sorting and clinics' capacity to accept new patients may affect observed patient flows. In addition, incoming patients' prescribing histories provide a noisy measure of the treatment style from which patients originate, potentially attenuating the OLS estimates toward zero. I therefore turn to closure-induced patient reallocations to obtain plausibly exogenous variation in the prescribing information received by clinics.

Table 4.2 presents the impact of receiving previously prescribed patients on prescribing to existing patients, estimated by instrumenting the potentially endogenous patient volume with

Table 4.2: Estimates of information effect on antibiotic prescribing intensity

	(1) IV	(2) IV
Number of prescribed patients	0.023*	0.023*
	(0.009)	(0.009)
Clinic covariates	No	Yes
Patient covariates	No	Yes
Clinic FE	Yes	Yes
Year-quarter FE	Yes	Yes
Observations	32,197	32,197
Outcome mean	-2.36	-2.36
First-stage F-stat	531.01	543.18

Notes: Number of prescribed patients and the outcome, antibiotic prescriptions per consultation, are in log-scale. Clinic covariates include clinic size, average physician age, share of female physicians, and share of physicians with an immigration background. Patient covariates include average patient age, share of children (age <18), share of female patients, and share of patients with an immigration background. Standard errors in parentheses are clustered at the receiving clinic level.

exposure to the practice styles of closed clinics. Column 1 indicates that, controlling only for clinic and time fixed effects, receiving twice as many patients with prior antibiotic prescriptions increases antibiotic prescribing intensity for existing patients by 2.3 percent. Controlling for clinic and patient covariates yields similar results. The increase in prescribing intensity to existing patients after receiving patients with a history of antibiotic treatment suggests that physicians may learn from peer physicians' practice styles through information passed along by moving patients. Such learning does not need to be efficiency-improving. In the case of antibiotic prescribing, where inefficiencies are often caused by over-treatment, high prescribing practice styles may be transmitted through a so far under-explored channel, moving patients.

4.2 Effect on prescribing quality

The baseline results show that exposure to incoming patients with prior prescriptions increases overall prescribing among existing patients. I next examine whether this response extends to the type of antibiotics prescribed and to the treatment process. Specifically, I consider broad- and narrow-spectrum prescribing together with diagnostic testing, prescriptions without prior testing, and follow-up prescriptions.

Table 4.3 summarizes the effects of patient information on prescribing intensity of antibiotics, classified according to their spectrum of activity. Columns (1) and (3) show the IV estimates for broad- and narrow-spectrum antibiotic prescribing intensity, including only clinic and time fixed effects. The estimates indicate that, when clinics receive double the number of prescribed patients, the prescribing intensity of broad-spectrum antibiotics for existing patients increases by 4.3 percent, which is almost twice the effect on narrow-spectrum antibiotic prescribing (2.2 percent). Controlling for clinic and patient covariates, doubling the incoming patients with

Table 4.3: Estimates of information effect on prescribing intensity by antibiotic class

	Broad-spectrum		Narrow-spectrum	
	(1) IV	(2) IV	(3) IV	(4) IV
Number of prescribed patients	0.043**	0.040*	0.022*	0.022*
	(0.015)	(0.016)	(0.011)	(0.011)
Clinic covariates	No	Yes	No	Yes
Patient covariates	No	Yes	No	Yes
Clinic FE	Yes	Yes	Yes	Yes
Year-quarter FE	Yes	Yes	Yes	Yes
Observations	32,197	32,197	32,197	32,197
Outcome mean	-3.93	-3.93	-2.83	-2.83
First-stage F-stat	531.01	543.18	531.01	543.18

Notes: Number of prescribed patients and the outcome, antibiotic prescriptions per consultation, are in log-scale. Clinic covariates include clinic size, average physician age, share of female physicians, and share of physicians with an immigration background. Patient covariates include average patient age, share of children (age <18), share of female patients, and share of patients with an immigration background. Antibiotics are classified into broad- and narrow-spectrum usage at ATC level 4 or 6. Standard errors in parentheses are clustered at the receiving clinic level.

any prior prescriptions corresponds to a 4 percent increase in the broad-spectrum antibiotic prescribing intensity among non-switchers, and the effect on narrow-spectrum prescribing remains unchanged. Broad-spectrum prescribing has a substantially lower outcome mean than narrow-spectrum prescribing, yet it responds more strongly to the patient inflow shock. This finding suggests that the transmitted practice style from origin clinics is especially salient on the more discretionary margin of prescribing. I find similar prescribing patterns using the share of prescribed patients as the strength of the information signal in Appendix Table E.2, where an additional 1 percentage point (0.01) in the share increases the broad- and narrow-spectrum prescribing intensity by 1.19 percent and 0.66 percent respectively.

The increase in narrow-spectrum prescribing suggests that spillovers do not operate solely through substitution into lower-quality treatment. Rather, these estimates indicate that practice styles from origin clinics spill over into the receiving clinic's overall treatment behavior. The finding is consistent with a mechanism of information transmission: once a patient enters the receiving clinic with an antibiotic treatment history, the threshold for treatment in the same drug class for existing patients may fall because the realized treatment reduces diagnostic uncertainty. This points to a more general shift in prescribing behavior rather than quality reduction or displacement of care for the incumbent patients alone. At the same time, the stronger increase in broad-spectrum prescribing highlights that the lower-quality margin of antibiotic use is sensitive to external information, and that the transmitted information may not necessarily be welfare-improving. In other words, patient inflows can diffuse both better and worse clinical practices. Exposure to prior treatment histories may improve treatment decisions on appropriate margins such as narrow-spectrum use, through information that may reinforce better practice styles and




Figure 4.1: IV estimates of information effect on prescribing quality indicators

Notes: The graph plots the estimates with 95% confidence intervals from regressing antibiotic prescribing intensity and three prescribing quality indicators respectively on the log-number of incoming patients with prior antibiotic prescriptions, instrumented with an IV computed from share-weighted mean prescribing at closed origin clinics. All outcomes are measured per-consultation and in log-scale for the existing patients. All regressions include the same set of clinic and patient covariates, as well as clinic and year-quarter fixed effects, as in the main specification for prescribing intensity. Standard errors are clustered at the receiving clinic level. Complete estimation results are in Appendix Table E.16.

allow for the discovery of new treatments, but it may also propagate lower-quality prescribing styles, especially if it crowds out other treatment tools, such as diagnostic tests and guidelines.

The increase in broad-spectrum prescribing suggests that information exposure may affect prescribing quality as well as overall prescribing intensity. To assess this interpretation, I next examine whether the response is accompanied by changes in diagnostic effort and treatment adjustment, using diagnostic testing, prescriptions without prior testing, and follow-up prescriptions as complementary outcomes.

Figure 4.1 presents the estimates for the information effect on these prescribing quality indicators alongside the estimate for overall prescribing intensity from Column (2) of Table 4.2, among existing patients at the receiving clinic. The estimated coefficients are positive for all four outcomes. The effects for all prescriptions, diagnostic tests, and prescriptions before testing are similar in magnitude and statistically significant. The coefficient for prescriptions with follow-up is also positive, although it is estimated less precisely.

The increase in overall prescribing is accompanied by an increase in diagnostic testing, indicating that clinics do not respond solely by expanding treatment while reducing clinical effort. One possible explanation is that patient inflows transmit not only prescribing outcomes as reflected by realized prescriptions, but also broader procedural practices from origin clinics. As a result, receiving clinics may change not just how much they prescribe, but also how often they use tests when treating conditions that may require antibiotics. Another possibility is that physicians differ in how they respond to the information carried by incoming patients. For physicians who remain hesitant to prescribe immediately, that information may primarily increase diagnostic testing rather than treatment itself, as prior treatment histories raise the salience of infection without fully displacing the demand for confirmation through timely testing. Under

Table 4.4: Estimates of information effect on prescribing intensity by antibiotic class

	(1) All prescriptions	(2) Broad-spectrum	(3) Narrow-spectrum
Number of prescribed patients	0.058**	0.060	0.070**
	(0.020)	(0.032)	(0.022)
Years since entry	0.004	-0.008	0.006
	(0.004)	(0.007)	(0.005)
Number of prescribed patients ×Years since entry	-0.002*	-0.001	-0.002**
	(0.001)	(0.001)	(0.001)
Clinic covariates	Yes	Yes	Yes
Patient covariates	Yes	Yes	Yes
Clinic FE	Yes	Yes	Yes
Year-quarter FE	Yes	Yes	Yes
Observations	32,197	32,197	32,197
Outcome mean	-2.36	-3.93	-2.83
First-stage F-stat	121.16	121.16	121.16

Notes: Number of prescribed patients and the outcome, antibiotic prescriptions per consultation, are in log-scale. Clinic covariates include clinic size, average physician age, share of female physicians, and share of physicians with an immigration background. Patient covariates include average patient age, share of children (age <18), share of female patients, and share of patients with an immigration background. Antibiotics are classified into broad- and narrow-spectrum usage at ATC level 4 or 6. Standard errors in parentheses are clustered at the receiving clinic level.

this interpretation, the informational spillover may be particularly strong on the testing margin.

On the other hand, while prescriptions issued before testing also increase, prescriptions with follow-up do not rise significantly. If follow-up prescriptions are interpreted as an indicator of lower initial prescribing quality, this finding suggests that the increase in prescribing without testing does not translate into a clear increase in subsequent corrective treatment. One interpretation is that physicians become more willing to prescribe prior to testing when they can draw on information embedded in patients' prior treatment histories. In that case, information transmitted through patient inflows may partly substitute for contemporaneous testing, allowing earlier prescribing without a detectable deterioration in this downstream quality measure. Overall, the results suggest that receiving clinics respond to the information brought in by incoming patients, and that the response is reflected not only in prescribing outcomes among existing patients, but also in treatment-related decisions such as diagnostic testing.

4.3 Heterogeneous response by clinic age

The baseline results identify an average response, but clinics may differ in how strongly they learn from the incoming information. In particular, belief updating upon the arrival of new information may vary with a clinic's years of operation, as newer clinics have less internal information to

Table 4.5: Estimates of information effect on prescribing quality indicators

	(1) All prescriptions	(2) Diagnostic tests	(3) Prescriptions before testing	(4) Prescriptions with follow-up
Number of prescribed patients	0.058**	0.051*	0.047*	0.063
	(0.020)	(0.022)	(0.021)	(0.036)
Years since entry	0.004	0.004	0.001	0.010
	(0.004)	(0.005)	(0.004)	(0.009)
Number of prescribed patients × Years since entry	-0.002*	-0.001	-0.001	-0.002
	(0.001)	(0.001)	(0.001)	(0.001)
Clinic covariates	Yes	Yes	Yes	Yes
Patient covariates	Yes	Yes	Yes	Yes
Clinic FE	Yes	Yes	Yes	Yes
Year-quarter FE	Yes	Yes	Yes	Yes
Observations	32,197	32,197	32,197	32,197
Outcome mean	-2.36	-2.20	-2.72	-4.97
First-stage F-stat	121.16	121.16	121.16	121.16

Notes: Number of prescribed patients and outcome are in log-scale. The outcome in column (1) is antibiotic prescriptions per consultation. Clinic covariates include clinic size, average physician age, share of female physicians, and share of physicians with an immigration background. Patient covariates include average patient age, share of children (age <18), share of female patients, and share of patients with an immigration background. Standard errors in parentheses are clustered at the receiving clinic level.

draw on, hence may be more responsive than clinics with more established practice routines.[19] To understand whether the information effect differs for newer and older clinics, I estimate an experience-specific effect by interacting the information measure with clinic age in years, calculated as the difference between the observed year and the year when the clinic first opened, with the following model:

$$y_{it} = \beta_1 info_{it} + \beta_2 info_{it} \times age_{it} + \beta_3 age_{it} + \delta X_{it} + FE_i + FE_t + \epsilon_{it}$$

Variable age_{it} measures a clinic i's years of operation in the market until time t. To obtain a causal estimate, I also interact the instrumental variable with age_{it}.

Column (1) of Table 4.4 indicates that every additional year of experience reduces the information effect from receiving patients with prior prescriptions by 0.2 percent. According to this estimate, when presented with twice the volume of information, a newly opened clinic within a year ($age_{it} = 0$) responds by prescribing 5.8 percent more upon a consultation from existing patients, whereas a 10-year-old clinic increases its prescribing intensity by 3.8 percent, 2 percentage points lower than the new clinic. This finding implies that exposure to incoming patients from prescribing-intensive origin clinics increases antibiotic prescribing among incumbent patients, but the size of this response declines as the receiving clinic becomes more established. In other words, younger clinics are more responsive to the information carried by incoming patients, while older clinics react less strongly, consistent with predictions from Bayesian belief updating models.

Similarly, for narrow-spectrum prescribing intensity, the positive main effect combined with the negative interaction implies that the impact of incoming information is largest when clinic age is low and becomes smaller with each additional year since entry. The estimates for broad-spectrum antibiotics show similar patterns, although not statistically significant. Given that broad-spectrum prescribing is much less common, as reflected in its lower outcome mean, the estimates are also less precise. The estimates also imply that the larger average effect on broad-spectrum prescribing in the baseline results (Table 4.3) is not primarily driven by differences across clinic age. Together with the result on overall antibiotic prescribing, I find that younger clinics respond to the external information more strongly, by shifting towards a more desirable category of antibiotic use.

Other quality indicators, such as the intensity of diagnostic testing and prescriptions without testing, show a similar pattern in Table 4.5. Younger clinics increase both testing and pre-testing prescribing among existing patients when exposed to incoming information, yet the absence of a significant interaction suggests weaker evidence that this response declines systematically with clinic age. As with the overall results from Figure 4.1, I find no statistically significant evidence of an information effect on prescriptions that require a corrective follow-up prescription.

In Appendix Tables E.12 and E.13, I show that the response to information does not vary

[19] Under a Bayesian learning framework, a clinic's years of operation may proxy for the precision of the priors because accumulated experience from observed patient outcomes can reduce uncertainty in diagnosis and treatment, thus making prior beliefs more precise as shown in Coscelli and Shum (2004). As priors become more precise, agents place less weight on new signals, implying weaker updating in response to the same information.

Table 4.6: Estimates of information effect on prescribing intensity using different windows

	(1) Baseline	(2) Past 4 quarters	(3) Past 8 quarters
Number of prescribed patients	0.023*		
	(0.009)		
Patients prescribed 4 quarters before switch		0.024*	
		(0.010)	
Patients prescribed 8 quarters before switch			0.023*
			(0.009)
Clinic covariates	Yes	Yes	Yes
Patient covariates	Yes	Yes	Yes
Clinic FE	Yes	Yes	Yes
Year-quarter FE	Yes	Yes	Yes
Observations	32,197	32,197	32,197
Outcome mean	-2.36	-2.36	-2.36
First-stage F-stat	530.41	514.94	537.44

Notes: Number of prescribed patients and the outcome, antibiotic prescriptions per consultation, are in log-scale. Clinic covariates include clinic size, average physician age, share of female physicians, and share of physicians with an immigration background. Patient covariates include average patient age, share of children (age <18), share of female patients, and share of patients with an immigration background. Standard errors in parentheses are clustered at the receiving clinic level.

by physician age. Therefore, within-clinic information stock seems to be more relevant than physician experience. Younger clinics are more sensitive to external information because they may have less settled workflows and weaker internal norms, which allow them to react more flexibly when they receive patients from origin clinics with different prior prescribing practices. Newer clinics that have fewer listed patients may also be more likely to exert effort in adjusting their treatment behavior to retain the existing patients. Older clinics, in contrast, are more likely to have established ways of treating patients, so the same incoming information has less effect on how they treat incumbents.

4.4 Robustness

When constructing the information signal, I use all switching patients between 2014 and 2019 who have received *any* antibiotic prescriptions, assuming that physicians at the receiving clinics view historical prescriptions as a per-patient uniform signal regardless of the prescribed date or volume. In Appendix Tables E.1 and E.4, I show that the estimated information effect is robust to alternative forms of this signal specification, including the share of prescribed patients and the raw number of prescribed patients instead of the log-number.

I further test the robustness of the signal specification by restricting the prescription date of the incoming patient, first because patients may be more likely to communicate recent prescriptions prior to switching, sending a stronger signal to the physicians, and physicians may

interpret newer prescriptions as a signal for recent developments or trends in the local prescribing environment. To ensure that the results are not driven by historical signals that may no longer be relevant, I employ alternative windows for prescriptions at the origin clinics of the switching patients. I define antibiotic prescriptions within a year or two (4 or 8 quarters) as the information that switching patients carry to the receiving clinics. Appendix Figure D.1 presents the distributions of the number of patients using varying information windows. The estimate in column (1) of Table 4.6 indicates that doubling the number of patients who received antibiotics in the past 4 quarters drives up the prescribing intensity to existing patients by 2.4 percent, 0.1 percentage point larger than the effect of the information signal constructed using a larger window of 8 quarters, and the effect from using all historical prescriptions. Although the magnitudes of the estimated effects are quite close, the result demonstrates the importance of newer prescribing signals.

As an additional check, I estimate the information effect on prescribing intensity, accounting for the variations in per-patient prescriptions. For each receiving clinic, I define the per-patient prescribing rate among incoming patients as the average number of prescriptions they received from their old clinics. This measure allows for varying weights of patients in the information signal composition and heterogeneous signal intensities for the receiving clinics, especially in cases where clinics receive the same number of prescribed patients who bring different volumes of prior prescriptions. This rate also reflects the past demand for antibiotics among the incoming patients, as well as the "ease" of obtaining an antibiotic prescription experienced at their origin clinics, which may shape their expectations for their new clinics. Patients with varying historical prescriptions may also actively communicate their needs and expectations, constituting heterogeneous signal strengths for the receiving clinics. Appendix Table E.7 presents the results using the per-patient prescribing rate as the information signal. Demonstrating that the coefficients remain stable across these varied specifications reinforces the argument that physicians are engaged in a deliberate process of belief revision based on the informational content of their caseload, rather than reacting to unobserved clinic-level shocks or simple changes in patient flow.

I also evaluate the sensitivity of the estimates to the instrument construction. The instrument linearly combines two components: one part represents the shock component, leveraging patients' redistribution after the as-good-as-random clinic closure; the other part proxies for clinic practice styles that influence the prescribing information these redistributed patients carry. In the main analysis, I use the average prescribing intensity of the closed clinics to non-switchers to proxy for the underlying practice style. While this average measure over all observed periods of a clinic until closure does predict the actual prescriptions, as is shown by the first stage regression of the instrument, it may not reflect the potential changes in prescribing styles over time. Prescribing styles may vary when physicians join or leave the team, or when factors such as information exposure gradually shift the prescribing behavior. Therefore, practice styles during a relatively shorter window may better capture the practice styles patients experience before closure. I use a moving average of prescribing intensity within 4 quarters as an alternative to probe the robustness of the instrument. To reduce the bias from changes induced by patient

movements, the prescribing intensity is again averaged over the non-switching patients. The estimated effects in Appendix Table E.10 are very similar to the estimates from the main specification. In fact, practice styles under the two definitions are distributed similarly in Appendix Figure D.2, which may be due to the relatively shorter span of the sample and the overall inertia in prescribing patterns.

4.5 Mechanism: information spillover or crowding out?

The model aims to estimate the effect of exposure to information brought by newly arrived patients. However, an overall increase in the number of listed patients may generate behavioral changes that could shift the quality of care delivered to the incumbent patients. With new patients coming in, physicians may be under increased time pressure and spend less time on each patient, and hence be more likely to prescribe without testing. Appointments may become harder to obtain, reducing the number of consultations for existing patients. In Appendix Table F.2, I examine the possibility of care resource crowding out by regressing the outcomes that can proxy for care utilization of the existing patients on the number of new patients from closed clinics, and show that receiving more patients does not affect the antibiotic prescribing intensity and consultations for existing patients. I only find a minor negative effect of -0.002 on the (log-)number of claims, indicating a 0.2 percent decrease in the number of claims filed for the incumbents when the patient intake doubles.

Additionally, Appendix Table F.1 summarizes the estimates from regressing the instrument on the same set of outcomes, demonstrating that the instrument, the proxy for information exposure, only has an impact on the prescribing intensity rather than the total number of consultations or claims. These findings provide little evidence that exposure to higher-prescribing origin clinics reduces service use for incumbent patients. Instead, the results suggest that the identifying variation is linked specifically to prescribing behavior rather than to a shift in service provision from existing to new patients.

Another falsification exercise lends credibility to the information mechanism. I relate the prescribing intensity of other drug classes to the instrument and the information measures for antibiotic prescribing. The candidate drugs are statins and drugs used in diabetes, two major groups of medications used to manage cardiovascular and metabolic diseases.[20] In primary care, they are typically used for long-term management of chronic diseases, whereas antibiotics are prescribed for short-term treatment of acute infections. Statins are prescribed to lower cholesterol and reduce cardiovascular risk, often taken daily for years or lifelong as part of prevention strategies. Similarly, A10 drugs are used to control blood glucose in diabetes, requiring ongoing monitoring and adherence over long periods. Therefore, an increasing number of patients per primary care provider can reduce the quality of chronic care because chronic disease management requires time-intensive, continuous monitoring and individualized follow-up.

Appendix Tables F.3 and F.4 show the results of regressing statin prescribing intensity to non-

[20] Statins are a class of cholesterol-lowering drugs under ATC level 4 class C10AA, HMG CoA reductase inhibitors. Drugs used in diabetes include all drugs under ATC class A10, including metformin, sulfonylureas, insulin, and newer agents like SGLT2 inhibitors or GLP-1 receptor agonists.

switchers on the instrument for exposure to information on antibiotics, and the two information measures. All three regressions yield null effects, supporting the hypothesis that the increase in prescribing is matched to the exposure to prescribing styles specific to antibiotics.

Several other pieces of evidence are also consistent with Bayesian learning. In Section 4.3, I show that responses to the information signal are larger in relatively new clinics, which are likely to have less precise priors under diagnostic uncertainty. At the same time, I find no heterogeneous effects by physician age, suggesting that within-clinic prescribing norms may matter more for prescribing decisions. I further examine heterogeneity by clinic structure in Appendix Table E.14, splitting the sample by the number of physicians in a clinic, and find that single-physician clinics' prescribing intensity increases more when exposed to incoming patients with prior prescriptions, while multi-physician clinics barely respond. One interpretation is that in a multi-physician clinic, physicians may learn not only from their own experience but also from colleagues through discussion, observation, and shared routines. This internal information exchange can make the clinic's prior more precise, so external information brought by incoming patients receives less weight. In a single-physician clinic, this internal source of information is weaker, leading to larger updating when new information arrives. Together, these findings also highlight the importance of the internal information environment of clinics in shaping medical decision-making.

5 Conclusion

Primary care physicians routinely make treatment decisions under uncertainty, relying on prior experience, accumulated knowledge, and imperfect signals from past cases. Yet relatively little is known about how these beliefs are formed and updated in practice, or how they translate into persistent prescribing styles. In this paper, I show that medical information, embedded in patients' prior antibiotic prescribing histories, can be transmitted across physicians through patient mobility. Exploiting quasi-random patient redistribution following primary care clinic closures in Denmark, I find that clinics receiving twice as many new patients with prior antibiotic prescriptions increase antibiotic prescribing intensity for their existing patients by 2.3 percent.

However, the practice styles that diffuse are not unambiguously welfare-improving. The increase is in both broad- and narrow-spectrum antibiotic prescribing, as well as in prescriptions issued before diagnostic testing. This pattern suggests that what spreads is not a narrowly defined low-quality practice, such as excessive broad-spectrum use alone, but a broader tendency toward more intensive antibiotic treatment. Meanwhile, receiving physicians also become more likely to conduct diagnostic tests, without a corresponding increase in follow-up prescriptions to correct the initial treatment. These results suggest that information from incoming patients' treatment histories can shift both prescribing decisions and related clinical procedures.

The heterogeneous responses across clinics further support an information-based interpretation. Newer clinics and single-physician clinics respond more strongly to prescribing signals, consistent with greater reliance on external information when internal experience is more limited. This pattern is in line with models of learning under uncertainty, in which agents place

greater weight on outside signals when their priors are less precise. It also helps distinguish information transmission from alternative mechanisms such as patient congestion, which would not necessarily predict stronger responses among clinics with less accumulated experience.

These findings contribute to the literature on physician learning and practice variation by identifying a novel information channel operating through patient mobility rather than direct peer interaction or institutional guidelines. The results suggest that physicians infer treatment norms from the prescribing histories of incoming patients and incorporate this information into their own decision-making. Importantly, this learning mechanism operates without direct interaction between physicians. Patients can act as carriers of medical information across providers, allowing clinic-specific practice styles to diffuse through movements within the healthcare system.

From a policy perspective, these findings have implications for the diffusion of prescribing practices and the design of interventions aimed at improving antibiotic stewardship and reducing unwarranted variation in care. Patient mobility can amplify both desirable and undesirable practices, implying that policies targeting individual physicians may generate spillover effects beyond the directly treated clinics. More broadly, the results show that informal information transmission can shape provider behavior even in a highly regulated primary care system with formal prescribing guidelines. Accounting for these spillovers is therefore important when evaluating the persistence of practice variation and the effectiveness of information-based interventions in healthcare markets.

References

Agha, L., Ericson, K. M., Geissler, K. H., and Rebitzer, J. B. (2022). Team Relationships and Performance: Evidence from Healthcare Referral Networks. *Management Science*, 68(5):3735–3754.

Agha, L. and Molitor, D. (2018). The Local Influence of Pioneer Investigators on Technology Adoption: Evidence from New Cancer Drugs. *The Review of Economics and Statistics*, 100(1):29–44.

Arrow, K. J., Bilir, L. K., and Sorensen, A. (2020). The Impact of Information Technology on the Diffusion of New Pharmaceuticals. *American Economic Journal: Applied Economics*, 12(3):1–39.

Avdic, D., Lundborg, P., and Vikström, J. (2019). Estimating returns to hospital volume: Evidence from advanced cancer surgery. *Journal of Health Economics*, 63:81–99.

Bailey, M., Johnston, D., Kuchler, T., Stroebel, J., and Wong, A. (2022). Peer Effects in Product Adoption. *American Economic Journal: Applied Economics*, 14(3):488–526.

Bennett, D., Hung, C.-L., and Lauderdale, T.-L. (2015). Health care competition and antibiotic use in Taiwan. *Journal of Industrial Economics*, 63(2):371–393.

Bhuller, M. and Sigstad, H. (2025). Feedback and Learning: The Causal Effects of Reversals on Judicial Decision-Making. *The Review of Economic Studies*, 92(4):2359–2397.

Chandra, A., Cutler, D., and Song, Z. (2011). Chapter Six - Who Ordered That? The Economics of Treatment Choices in Medical Care. In Pauly, M. V., Mcguire, T. G., and Barros, P. P., editors, *Handbook of Health Economics*, volume 2, pages 397–432.

Chandra, A. and Staiger, D. O. (2007). Productivity spillovers in health care: evidence from the treatment of heart attacks. *Journal of Political Economy*, 115(1):103–140.

Chen, Y. (2021). Team-Specific Human Capital and Team Performance: Evidence from Doctors. *American Economic Review*, 111(12):3923–3962.

Ching, A. T. (2010). Consumer learning and heterogeneity: Dynamics of demand for prescription drugs after patent expiration. *International Journal of Industrial Organization*, 28(6):619–638.

Coleman, J., Katz, E., and Menzel, H. (1957). The Diffusion of an Innovation Among Physicians. *Sociometry*, 20(4):253–270.

Coscelli, A. and Shum, M. (2004). An empirical model of learning and patient spillovers in new drug entry. *Journal of Econometrics*, 122(2):213–246.

Crawford, G. S. and Shum, M. (2005). Uncertainty and Learning in Pharmaceutical Demand. *Econometrica*, 73(4):1137–1173.

Currie, J., Lin, W., and Meng, J. (2014). Addressing antibiotic abuse in China: An experimental audit study. *Journal of Development Economics*, 110:39–51.

Currie, J. M. and MacLeod, W. B. (2020). Understanding Doctor Decision Making: The Case of Depression Treatment. *Econometrica*, 88(3):847–878.

Cutler, D., Skinner, J. S., Stern, A. D., and Wennberg, D. (2019). Physician Beliefs and Patient Preferences: A New Look at Regional Variation in Health Care Spending. *American Economic Journal: Economic Policy*, 11(1):192–221.

Domenella, D. (2025). Market Size and the Returns to Surgeon Volume: Evidence from Joint Replacements. Job Market Paper.

Epstein, A. J. and Ketcham, J. D. (2014). Information technology and agency in physicians' prescribing decisions. *The RAND Journal of Economics*, 45(2):422–448.

Epstein, A. J. and Nicholson, S. (2009). The formation and evolution of physician treatment styles: an application to cesarean sections. *Journal of Health Economics*, 28(6):1126–1140.

Fadlon, I. and Van Parys, J. (2020). Primary care physician practice styles and patient care: Evidence from physician exits in medicare. *Journal of Health Economics*, 71:102304.

Finkelstein, A., Gentzkow, M., and Williams, H. (2016). Sources of geographic variation in health care: Evidence from patient migration. *Quarterly Journal of Economics*, 131(4):1681–1726.

Gaynor, M., Seider, H., and Vogt, W. B. (2005). The Volume-Outcome Effect, Scale Economies, and Learning-by-Doing. *American Economic Review*, 95(2):243–247.

Haastrup, P., Møller, A., Kristensen, J. K., and Huibers, L. (2025). Danish primary care: A focus on general practice in the Danish healthcare system. *Scandinavian Journal of Primary Health Care*, 43(4):839–845.

Huang, S. and Ullrich, H. (2023). Provider effects in antibiotic prescribing: Evidence from physician exits. Berlin School of Economics Discussion Papers Nr. 18.

Huang, S. and Ullrich, H. (2024). Provider effects in antibiotic prescribing: Evidence from physician exits. *Journal of Human Resources*.

Kjaersgaard, M. I. S., Vedsted, P., Parner, E. T., Bech, B. H., Vestergaard, M., Flarup, K. R., and Fenger-Grøn, M. (2016). Algorithm linking patients and general practices in Denmark using the Danish National Health Service Register. *Clinical Epidemiology*, 8:273–283.

Laxminarayan, R., Duse, A., Wattal, C., Zaidi, A. K. M., Wertheim, H. F. L., Sumpradit, N., Vlieghe, E., Hara, G. L., Gould, I. M., Goossens, H., Greko, C., So, A. D., Bigdeli, M., Tomson, G., Woodhouse, W., Ombaka, E., Peralta, A. Q., Qamar, F. N., Mir, F., Kariuki, S., Bhutta, Z. A., Coates, A., Bergstrom, R., Wright, G. D., Brown, E. D., and Cars, O. (2013). Antibiotic resistance – the need for global solutions. *The Lancet Infectious Diseases Commission*, 13(12):1057–1098.

Manski, C. F. (2018). Reasonable patient care under uncertainty. *Health Economics*, 27(10):1397–1421.

Molitor, D. (2018). The evolution of physician practice styles: evidence from cardiologist migration. *American Economic Journal: Economic Policy*, 10(1):326–56.

Phelps, C. E. (1992). Diffusion of Information in Medical Care. *Journal of Economic Perspectives*, 6(3):23–42.

Phelps, C. E. (2000). Information diffusion and best practice adoption. In *Handbook of health economics*, volume 1, pages 223–264. Elsevier.

Ribers, M. A. and Ullrich, H. (2023). Machine learning and physician prescribing: a path to reduced antibiotic use. Berlin School of Economics Discussion Paper Nr. 19.

Rockoff, J. E., Staiger, D. O., Kane, T. J., and Taylor, E. S. (2012). Information and Employee Evaluation: Evidence from a Randomized Intervention in Public Schools. *American Economic Review*, 102(7):3184–3213.

Simonsen, M., Skipper, L., Skipper, N., and Thingholm, P. R. (2021). Discontinuity in care: Practice closures among primary care providers and patient health care utilization. *Journal of Health Economics*, 80:102551.

Sorensen, A. T. (2006). Social learning and health plan choice. *The Rand Journal of Economics*, 37(4):1–29.

Zhu, Z. (2023). The Value of Patients: Heterogenous Physician Learning and Generic Drug Diffusion. Working Paper.

A Antibiotic classifications

Table A.1: Classification of antibiotics


ATC4 or ATC6 class	Name
Broad-spectrum Antibiotics	
J01CR	Combinations of penicillins, including beta-lactamase inhibitors
J01DC	Second-generation cephalosporins
J01DD	Third-generation cephalosporins
J01(FA-FA01)	Macrolides except erythromycin
J01MA	Fluoroquinolones
Narrow-spectrum Antibiotics	
J01CA	Penicillins with extended spectrum
J01CE	Beta-lactamase-sensitive penicillins
J01CF	Beta-lactamase-resistant penicillins
J01DB	First-generation cephalosporins
J01FA01	Erythromycin
Other Antibiotics	
J01EA	Trimethoprim and derivatives
J01EB	Short-acting sulfonamides
J01EE	Combinations of sulfonamides and trimethoprim, including derivatives
J01XE	Nitrofuran derivatives
J01XX	Other antibacterials

B Patient descriptive statistics


Table B.1: Patient descriptive statistics

	mean (s.d.)	
	Non-switchers	Switchers
Number of antibiotic prescriptions	1.32	1.29
	(0.79)	(0.72)
Number of consultations	1.94	1.91
	(1.53)	(1.44)
Number of claims	4.41	4.29
	(4.37)	(4.15)
Share of female patients	0.51	0.55
	(0.50)	(0.50)
Patient age	43.29	39.53
	(24.09)	(22.72)
Share of patients with immigration background	0.10	0.13
	(0.31)	(0.33)
Number of switches		1.34
		(0.67)
Age when switching		38.22
		(22.47)
Observations	121,250,996	

Note: This table summarizes the mean and the standard deviation of the listed variables from an unbalanced quarterly panel consisting of 5,011,537 patients assigned to 1,838 active primary care clinics from 2014 to 2019. Patients with an immigration background include patients who are first- or second-generation immigrants.



(a) Difference in home distance to new and original clinic



(b) Difference in prescribing style between new and original clinic

Notes: Distances are calculated from the WGS84 (World Geodetic System 1984) coordinates of pseudonymized residential and clinic addresses using the Haversine formula method. Differences in distances to the chosen clinics are calculated for each switch observed during the sample period, and trimmed at +/- 10km to ensure anonymity. Prescribing style of a clinic is defined as the mean antibiotic prescribing intensity among non-switching patients over all observations of a given clinic. Prescribing intensities are calculated from the number of antibiotic prescriptions over the number of consultations. Differences in prescribing styles are trimmed at +/- 0.2 to ensure anonymity.

C Pre-trend and balance tests

C.1 Pre-trend test for the instrument

Table C.1: Outcome: prescribing intensity

	(1) OLS	(2) OLS	(3) OLS
IV: Mean prescribing of closed clinics	0.032*		
	(0.013)		
IV, t+1		0.023	0.023
		(0.013)	(0.012)
IV, t+2			0.004
			(0.013)
Lagged clinic covariates	Yes	Yes	Yes
Lagged patient covariates	Yes	Yes	Yes
Clinic FE	Yes	Yes	Yes
Year-quarter FE	Yes	Yes	Yes
Observations	29,112	26,610	24,354
Outcome mean	-2.37	-2.38	-2.38
R-squared			

Notes: Number of prescribed patients and the outcome, antibiotic prescriptions per consultation, are in log-scale. Clinic covariates include clinic size, average physician age, share of female physicians, and share of physicians with an immigration background. Patient covariates include average patient age, share of children (age <18), share of female patients, and share of patients with an immigration background. Standard errors in parentheses are clustered at the receiving clinic level.

C.2 Balance test for the instrument




Figure C.1: Balance test for clinic activities among all patients

Notes: The graph plots the estimates with 95% confidence intervals from regressing the variables respectively on the lead instrument, controlling for time and clinic fixed effects. *Total number of patients*, *Claims per physician*, *Consultations*, *Prescribing intensity* are in log-scale. Standard errors are clustered at the receiving clinic level.




Figure C.2: Balance test for clinic activities among non-switchers

Notes: The graph plots the estimates with 95% confidence intervals from regressing the variables respectively on the lead instrument, controlling for time and clinic fixed effects. *Located in larger cities* is a 0/1 indicator for clinics located in Copenhagen, Aarhus, and Odense. *Claim per non-switching patient*, *Consultation per non-switching patient*, *Testing intensity*, and *Prescribing intensity* are in log-scale. Standard errors are clustered at the receiving clinic level.

Table C.2: Balance table for the instrument

	Coef.	Mean
Claim per non-switching patient	-.016	2.47
	(0.019)	
Claims per physician	.004	8.28
	(0.010)	
Clinic size	.013	2.23
	(0.021)	
Consultation per non-switching patient	-.002	.84
	(0.007)	
Consultations (all patients)	.015	7.82
	(0.009)	
Patient age	-.025	42.05
	(0.084)	
Patients with immigration background	-.001	.13
	(0.001)	
Physician age	.167	53.43
	(0.184)	
Physicians with immigration background	0	.08
	(0.004)	
Prescribing intensity (all patients)	.011	-2.31
	(0.009)	
Share of children	-.002	.22
	(0.001)	
Share of female patients	.001	.49
	(0.001)	
Share of female physicians	-.006	.49
	(0.008)	
Testing intensity	.016	-2.2
	(0.016)	
Total number of patients	-.01	7.89
	(0.007)	
Total quarters since entry	-.034	4.12
	(0.021)	

Notes: This table summarizes the estimates, standard errors (in parentheses), and outcome mean for regressing each of the variables on the lead instrument, calculated as the mean prescribing of closed clinics. The regressions control for time and clinic fixed effects, and cluster standard errors at the clinic level.

C.3 Balance test for clinic closure





Figure C.3: Estimates of the effects of selected variables on clinic closure


Notes: The graph plots the estimates with 95% confidence intervals from regressing a lagged 0/1 closure indicator jointly on the selected variables, controlling for time and clinic fixed effects. *Located in larger cities* is a 0/1 indicator for clinics located in Copenhagen, Aarhus, and Odense. *Prescribing intensity*, *Total revenue*, and *Claims per patient* are in log-scale. Standard errors are clustered at the receiving clinic level.

D Distribution of information measures and instruments

D.1 Distributions of patient volumes with different prescribing windows



(a) Patients prescribed within 4 quarters before switch



(b) Patients prescribed within 8 quarters before switch

Figure D.1: Distributions of incoming patients with prior prescriptions

Notes: These figures present the distributions of the information measure, defined as the number of incoming patients with prior prescribing histories of antibiotics, while restricting the prescription window to 4 and 8 quarters before switching. The number of patients prescribed anytime before switching serves as the main information measure. Number of patients is in log-scale. The log-number of patients is trimmed at 4.4 to ensure anonymity.

D.2 Distribution of the instrument using an alternative practice style definition




Figure D.2: Distribution of instruments

Notes: The graph plots the distributions and the mean values of the instrument constructed for the receiving clinics, using a time-invariant mean antibiotic prescribing to the non-switchers across the entire observational period as practice style, against the instrument constructed using a moving average of antibiotic prescribing intensity to the non-switchers within 4 quarters. The distribution is trimmed at an instrument value of 0.4 to ensure anonymity.

E Additional Results

E.1 Share of prescribed patients as information measure

The intensity of information exposure is measured as the share of incoming patients who were prescribed the treatment by their previous physicians, among all the new patients that the clinic received in a given quarter:

$$share_{it} = \frac{\sum_{j=1}^{J}(prescribed_{jt} \times switch_{ijt})}{\sum_{j=1}^{J} switch_{ijt}}.$$

Table E.1: Estimates of information effect on prescribing intensity

	(1) IV	(2) IV
Share of prescribed patients	0.69*	0.68*
	(0.29)	(0.29)
Clinic covariates	No	Yes
Patient covariates	No	Yes
Clinic FE	Yes	Yes
Year-quarter FE	Yes	Yes
Observations	32,197	32,197
Outcome mean	-2.36	-2.36
First-stage F-stat	48.60	47.95

Notes: The outcome, antibiotic prescriptions per consultation, is in log-scale. Clinic covariates include clinic size, average physician age, share of female physicians, and share of physicians with an immigration background. Patient covariates include average patient age, share of children (age <18), share of female patients, and share of patients with an immigration background. Standard errors in parentheses are clustered at the receiving clinic level.

Table E.2: Estimates of information effect on prescribing intensity by antibiotic class

	Broad-spectrum		Narrow-spectrum	
	(1)	(2)	(3)	(4)
	IV	IV	IV	IV
Share of prescribed patients	1.28**	1.19*	0.66*	0.66*
	(0.48)	(0.49)	(0.33)	(0.33)
Clinic covariates	No	Yes	No	Yes
Patient covariates	No	Yes	No	Yes
Clinic FE	Yes	Yes	Yes	Yes
Year-quarter FE	Yes	Yes	Yes	Yes
Observations	32,197	32,197	32,197	32,197
Outcome mean	-3.93	-3.93	-2.83	-2.83
First-stage F-stat	48.60	47.95	48.60	47.95

Notes: The outcome, antibiotic prescriptions per consultation, is in log-scale. Clinic covariates include clinic size, average physician age, share of female physicians, and share of physicians with an immigration background. Patient covariates include average patient age, share of children (age <18), share of female patients, and share of patients with an immigration background. Antibiotics are classified into broad- and narrow-spectrum usage at ATC level 4 or 6. Standard errors in parentheses are clustered at the receiving clinic level.

Table E.3: Estimates of information effect on prescribing quality indicators

	(1) All prescriptions	(2) Diagnostic tests	(3) Prescriptions before testing	(4) Prescriptions with follow-up
Share of prescribed patients	0.68*	0.73*	0.60	0.68
	(0.29)	(0.35)	(0.31)	(0.58)
Clinic covariates	Yes	Yes	Yes	Yes
Patient covariates	Yes	Yes	Yes	Yes
Clinic FE	Yes	Yes	Yes	Yes
Year-quarter FE	Yes	Yes	Yes	Yes
Observations	32,197	32,197	32,197	32,197
Outcome mean	-2.36	-2.20	-2.72	-4.97
First-stage F-stat	47.95	47.95	47.95	47.95

Notes: All outcomes are measured per-consultation and in log-scale; the outcome in column (1) is antibiotic prescriptions per consultation. Number of prescribed patients is in log-scale. Clinic covariates include clinic size, average physician age, share of female physicians, and share of physicians with an immigration background. Patient covariates include average patient age, share of children (age <18), share of female patients, and share of patients with an immigration background. Standard errors in parentheses are clustered at the receiving clinic level.

E.2 Count of prescribed patients as information measure

Table E.4: Estimates of information effect on prescribing intensity

	(1) IV	(2) IV
Number of prescribed patients	0.0018*	0.0018*
	(0.0007)	(0.0008)
Clinic covariates	No	Yes
Patient covariates	No	Yes
Clinic FE	Yes	Yes
Year-quarter FE	Yes	Yes
Observations	32,197	32,197
Outcome mean	-2.36	-2.36
First-stage F-stat	115.67	120.66

Notes: Number of prescribed patients is the raw count of patients with prior prescriptions. The outcome, antibiotic prescriptions per consultation, is in log-scale. Clinic covariates include clinic size, average physician age, share of female physicians, and share of physicians with an immigration background. Patient covariates include average patient age, share of children (age <18), share of female patients, and share of patients with an immigration background. Standard errors in parentheses are clustered at the receiving clinic level.

Table E.5: Estimates of information effect on prescribing intensity by antibiotic class

	Broad-spectrum		Narrow-spectrum	
	(1)	(2)	(3)	(4)
	IV	IV	IV	IV
Number of prescribed patients	0.003**	0.003*	0.002*	0.002*
	(0.001)	(0.001)	(0.001)	(0.001)
Clinic covariates	No	Yes	No	Yes
Patient covariates	No	Yes	No	Yes
Clinic FE	Yes	Yes	Yes	Yes
Year-quarter FE	Yes	Yes	Yes	Yes
Observations	32,197	32,197	32,197	32,197
Outcome mean	-3.93	-3.93	-2.83	-2.83
First-stage F-stat	115.67	120.66	115.67	120.66

Notes: Number of prescribed patients is the raw count of patients with prior prescriptions. The outcome, antibiotic prescriptions per consultation, is in log-scale. Clinic covariates include clinic size, average physician age, share of female physicians, and share of physicians with an immigration background. Patient covariates include average patient age, share of children (age <18), share of female patients, and share of patients with an immigration background. Antibiotics are classified into broad- and narrow-spectrum usage at ATC level 4 or 6. Standard errors in parentheses are clustered at the receiving clinic level.

Table E.6: Estimates of information effect on prescribing quality indicators

	(1) All prescriptions	(2) Diagnostic tests	(3) Prescriptions before testing	(4) Prescriptions with follow-up
Number of prescribed patients	0.002*	0.002*	0.002	0.002
	(0.001)	(0.001)	(0.001)	(0.002)
Clinic covariates	Yes	Yes	Yes	Yes
Patient covariates	Yes	Yes	Yes	Yes
Clinic FE	Yes	Yes	Yes	Yes
Year-quarter FE	Yes	Yes	Yes	Yes
Observations	32,197	32,197	32,197	32,197
Outcome mean	-2.36	-2.20	-2.72	-4.97
First-stage F-stat	120.66	120.66	120.66	120.66

Notes: Number of prescribed patients is the raw count of patients with prior prescriptions. All outcomes are measured per-consultation and in log-scale; the outcome in column (1) is antibiotic prescriptions per consultation. Clinic covariates include clinic size, average physician age, share of female physicians, and share of physicians with an immigration background. Patient covariates include average patient age, share of children (age <18), share of female patients, and share of patients with an immigration background. Standard errors in parentheses are clustered at the receiving clinic level.

E.3 Prescriptions per incoming patient as information measure

Table E.7: Estimates of information effect on prescribing intensity

	(1) IV	(2) IV
Prescriptions per new patient	0.17*	0.18*
	(0.07)	(0.08)
Clinic covariates	No	Yes
Patient covariates	No	Yes
Clinic FE	Yes	Yes
Year-quarter FE	Yes	Yes
Observations	32,197	32,197
Outcome mean	-2.36	-2.36
First-stage F-stat	48.88	46.38

Notes: Number of prescribed patients is the raw count of patients with prior prescriptions. The outcome, antibiotic prescriptions per consultation, is in log-scale. Clinic covariates include clinic size, average physician age, share of female physicians, and share of physicians with an immigration background. Patient covariates include average patient age, share of children (age <18), share of female patients, and share of patients with an immigration background. Standard errors in parentheses are clustered at the receiving clinic level.

I test the model using another definition of information that accounts for the variations in per-patient prescriptions. For each receiving clinic i, I define the per-patient prescribing rate among incoming patients as the average number of prescriptions they received from their old clinics. Formally:

$$rate_{it} = \frac{\sum_{j=1}^{J}(\#prescribed_{jkt} \times switch_{ijt})}{\sum_{j=1}^{J} switch_{ijt}}$$

where $\#prescribed_{jkt}$ is the total number of prescriptions patient j at their origin clinic before switching to destination clinic i at period t.

Table E.8: Estimates of information effect on prescribing intensity by antibiotic class

	Broad-spectrum		Narrow-spectrum	
	(1)	(2)	(3)	(4)
	IV	IV	IV	IV
Prescriptions per new patient	0.32**	0.31*	0.17*	0.17*
	(0.12)	(0.13)	(0.08)	(0.09)
Clinic covariates	No	Yes	No	Yes
Patient covariates	No	Yes	No	Yes
Clinic FE	Yes	Yes	Yes	Yes
Year-quarter FE	Yes	Yes	Yes	Yes
Observations	32,197	32,197	32,197	32,197
Outcome mean	-3.93	-3.93	-2.83	-2.83
First-stage F-stat	48.88	46.38	48.88	46.38

Notes: Number of prescribed patients is the raw count of patients with prior prescriptions. The outcome, antibiotic prescriptions per consultation, is in log-scale. Clinic covariates include clinic size, average physician age, share of female physicians, and share of physicians with an immigration background. Patient covariates include average patient age, share of children (age <18), share of female patients, and share of patients with an immigration background. Antibiotics are classified into broad- and narrow-spectrum usage at ATC level 4 or 6. Standard errors in parentheses are clustered at the receiving clinic level.

Table E.9: Estimates of information effect on prescribing quality indicators

	(1) All prescriptions	(2) Diagnostic tests	(3) Prescriptions before testing	(4) Prescriptions with follow-up
Prescriptions per new patient	0.18*	0.19*	0.15	0.18
	(0.08)	(0.09)	(0.08)	(0.15)
Clinic covariates	Yes	Yes	Yes	Yes
Patient covariates	Yes	Yes	Yes	Yes
Clinic FE	Yes	Yes	Yes	Yes
Year-quarter FE	Yes	Yes	Yes	Yes
Observations	32,197	32,197	32,197	32,197
Outcome mean	-2.36	-2.20	-2.72	-4.97
First-stage F-stat	46.38	46.38	46.38	46.38

Notes: Number of prescribed patients is the raw count of patients with prior prescriptions. All outcomes are measured per-consultation and in log-scale; the outcome in column (1) is antibiotic prescriptions per consultation. Clinic covariates include clinic size, average physician age, share of female physicians, and share of physicians with an immigration background. Patient covariates include average patient age, share of children (age <18), share of female patients, and share of patients with an immigration background. Standard errors in parentheses are clustered at the receiving clinic level.

E.4 Alternative practice style measure

Table E.10: Estimates of effects on prescribing intensity using moving average

	(1) IV	(2) IV
Number of prescribed patients	0.026**	0.026*
	(0.010)	(0.010)
Clinic covariates	No	Yes
Patient covariates	No	Yes
Clinic FE	Yes	Yes
Year-quarter FE	Yes	Yes
Observations	32,197	32,197
Outcome mean	-2.36	-2.36
First-stage F-stat	402.48	413.38

Notes: Number of prescribed patients and the outcome, antibiotic prescriptions per consultation, are in log-scale. Clinic covariates include clinic size, average physician age, share of female physicians, and share of physicians with an immigration background. Patient covariates include average patient age, share of children (age <18), share of female patients, and share of patients with an immigration background. Standard errors in parentheses are clustered at the receiving clinic level.

E.5 Adjustments to the clinic sample

Table E.11: Estimates of information effect on prescribing intensity

	(1) Excluding takeovers	(2) Excluding new clinics
Number of prescribed patients	0.020*	0.020*
	(0.009)	(0.009)
Clinic covariates	Yes	Yes
Patient covariates	Yes	Yes
Clinic FE	Yes	Yes
Year-quarter FE	Yes	Yes
Observations	30,384	28,117
Outcome mean	-2.35	-2.34
First-stage F-stat	503.72	451.47

Notes: Number of prescribed patients and the outcome, antibiotic prescriptions per consultation, are in log-scale. Clinic covariates include clinic size, average physician age, share of female physicians, and share of physicians with an immigration background. Patient covariates include average patient age, share of children (age <18), share of female patients, and share of patients with an immigration background. Standard errors in parentheses are clustered at the receiving clinic level.

E.6 Information interacted with physician age

Table E.12: Estimates of information effect interacted with physician age on prescribing intensity

	(1) All prescriptions	(2) Broad-spectrum	(3) Narrow-spectrum
Number of prescribed patients	0.012	0.107	0.029
	(0.063)	(0.102)	(0.075)
Physician age	0.002	0.011*	0.002
	(0.003)	(0.005)	(0.004)
Number of prescribed patients × Physician age	0.000	-0.001	-0.000
	(0.001)	(0.002)	(0.001)
Clinic covariates	Yes	Yes	Yes
Patient covariates	Yes	Yes	Yes
Clinic FE	Yes	Yes	Yes
Year-quarter FE	Yes	Yes	Yes
Observations	32,197	32,197	32,197
Outcome mean	-2.36	-3.93	-2.83
First-stage F-stat	116.01	116.01	116.01

Notes: Number of prescribed patients and the outcome, antibiotic prescriptions per consultation, are in log-scale. Clinic covariates include clinic size, average physician age, share of female physicians, and share of physicians with an immigration background. Patient covariates include average patient age, share of children (age <18), share of female patients, and share of patients with an immigration background. Antibiotics are classified into broad- and narrow-spectrum usage at ATC level 4 or 6. Standard errors in parentheses are clustered at the receiving clinic level.

Table E.13: Estimates of information effect interacted with physician age on prescribing quality indicators

	(1) All prescriptions	(2) Diagnostic tests	(3) Prescriptions before testing	(4) Prescriptions with follow-up
Number of prescribed patients	0.012	0.042	0.001	0.034
	(0.063)	(0.075)	(0.069)	(0.122)
Physician age	0.002	0.003	0.002	0.000
	(0.003)	(0.004)	(0.003)	(0.006)
Number of prescribed patients × Physician age	0.000	-0.000	0.000	-0.000
	(0.001)	(0.001)	(0.001)	(0.002)
Clinic covariates	Yes	Yes	Yes	Yes
Patient covariates	Yes	Yes	Yes	Yes
Clinic FE	Yes	Yes	Yes	Yes
Year-quarter FE	Yes	Yes	Yes	Yes
Observations	32,197	32,197	32,197	32,197
Outcome mean	-2.36	-2.20	-2.72	-4.97
First-stage F-stat	116.01	116.01	116.01	116.01

Notes: Number of prescribed patients and outcome are in log-scale. The outcome in column (1) is antibiotic prescriptions per consultation. Clinic covariates include clinic size, average physician age, share of female physicians, and share of physicians with an immigration background. Patient covariates include average patient age, share of children (age <18), share of female patients, and share of patients with an immigration background. Standard errors in parentheses are clustered at the receiving clinic level.

E.7 Information effect on single-physician clinics

Table E.14: Estimates of information effect on prescribing intensity

	(1) Single-physician	(2) Multiple-physician
Number of prescribed patients	0.047**	0.003
	(0.016)	(0.010)
Clinic covariates	Yes	Yes
Patient covariates	Yes	Yes
Clinic FE	Yes	Yes
Year-quarter FE	Yes	Yes
Observations	13,490	18,679
Outcome mean	-2.29	-2.41
First-stage F-stat	272.93	309.18

Notes: Number of prescribed patients and the outcome, antibiotic prescriptions per consultation, are in log-scale. Clinic covariates include clinic size, average physician age, share of female physicians, and share of physicians with an immigration background. Patient covariates include average patient age, share of children (age <18), share of female patients, and share of patients with an immigration background. Antibiotics are classified into broad- and narrow-spectrum usage at ATC level 4 or 6. Standard errors in parentheses are clustered at the receiving clinic level.

Table E.15: Estimates of information effect on prescribing quality indicators of single-physician clinics

	(1) All prescriptions	(2) Diagnostic tests	(3) Prescriptions before testing	(4) Prescriptions with follow-up
Number of prescribed patients	0.047**	0.047*	0.042*	0.032
	(0.016)	(0.020)	(0.017)	(0.032)
Clinic covariates	Yes	Yes	Yes	Yes
Patient covariates	Yes	Yes	Yes	Yes
Clinic FE	Yes	Yes	Yes	Yes
Year-quarter FE	Yes	Yes	Yes	Yes
Observations	13,490	13,490	13,490	13,490
Outcome mean	-2.29	-2.28	-2.62	-4.81
First-stage F-stat	272.93	272.93	272.93	272.93

Notes: Number of prescribed patients is the raw count of patients with prior prescriptions. All outcomes are measured per-consultation and in log-scale; the outcome in column (1) is antibiotic prescriptions per consultation. Clinic covariates include clinic size, average physician age, share of female physicians, and share of physicians with an immigration background. Patient covariates include average patient age, share of children (age <18), share of female patients, and share of patients with an immigration background. Standard errors in parentheses are clustered at the receiving clinic level.

E.8 Companion tables to graphs

Table E.16: Estimates of information effect on prescribing quality indicators

	(1) All prescriptions	(2) Diagnostic tests	(3) Prescriptions before testing	(4) Prescriptions with follow-up
Number of prescribed patients	0.023*	0.025*	0.020*	0.023
	(0.009)	(0.011)	(0.010)	(0.019)
Patient age	0.006	0.011**	0.003	-0.036***
	(0.003)	(0.004)	(0.004)	(0.005)
Share of children	0.207	0.181	0.159	-2.665***
	(0.239)	(0.298)	(0.262)	(0.350)
Share of female patients	0.523	0.884*	0.169	-0.816*
	(0.274)	(0.343)	(0.305)	(0.407)
Patients with immigration background	-0.027	0.114	0.002	-0.671*
	(0.221)	(0.263)	(0.244)	(0.298)
Physician age	0.002*	0.002*	0.003**	-0.000
	(0.001)	(0.001)	(0.001)	(0.002)
Share of female physicians	-0.026	0.049	-0.050	-0.033
	(0.022)	(0.032)	(0.027)	(0.038)
Clinic size	-0.007	-0.021**	-0.004	-0.018
	(0.006)	(0.008)	(0.007)	(0.011)
Physicians with immigration background	0.013	-0.018	0.019	0.018
	(0.036)	(0.040)	(0.039)	(0.068)
Clinic FE	Yes	Yes	Yes	Yes
Year-quarter FE	Yes	Yes	Yes	Yes
Observations	32,197	32,197	32,197	32,197
Outcome mean	-2.36	-2.20	-2.72	-4.97
First-stage F-stat	543.182	543.182	543.182	543.182

Notes: All outcomes are measured per-consultation and in log-scale; the outcome in column (1) is antibiotic prescriptions per consultation. Number of prescribed patients is in log-scale. Clinic covariates include clinic size, average physician age, share of female physicians, and share of physicians with an immigration background. Patient covariates include average patient age, share of children (age <18), share of female patients, and share of patients with an immigration background. Standard errors in parentheses are clustered at the receiving clinic level.

F Tests for crowding out

F.1 Care utilization outcomes

Table F.1: Reduced form OLS estimates of effects of the IV

	(1) Prescribing intensity	(2) Consultations	(3) Claims
Mean prescribing at origin clinic	0.032*	-0.012	-0.009
	(0.013)	(0.008)	(0.008)
Lagged clinic covariates	Yes	Yes	Yes
Lagged patient covariates	Yes	Yes	Yes
Clinic FE	Yes	Yes	Yes
Year-quarter FE	Yes	Yes	Yes
Observations	29,112	29,112	29,112
Outcome mean	-2.37	7.09	8.17
R-squared	0.78	0.99	0.99

Notes: All outcomes are in log-scale and measured for the existing patients (non-switchers); the prescribing intensity outcome is antibiotic prescriptions per consultation. Clinic covariates include clinic size, average physician age, share of female physicians, and share of physicians with an immigration background. Patient covariates include average patient age, share of children (age <18), share of female patients, and share of patients with an immigration background. Standard errors in parentheses are clustered at the receiving clinic level.

Table F.2: Reduced form OLS estimates of effects of the number of patients from closed clinics

	(1) Prescribing intensity	(2) Consultations	(3) Claims
Number of new patients	0.001	-0.002	-0.002*
	(0.001)	(0.001)	(0.001)
Lagged clinic covariates	Yes	Yes	Yes
Lagged patient covariates	Yes	Yes	Yes
Clinic FE	Yes	Yes	Yes
Year-quarter FE	Yes	Yes	Yes
Observations	29,112	29,112	29,112
Outcome mean	-2.37	7.09	8.17
R-squared	0.78	0.99	0.99

Notes: All outcomes are in log-scale and measured for the existing patients (non-switchers); the prescribing intensity outcome is antibiotic prescriptions per consultation. Number of new patients is in log-scale and refers to the number of patients coming from a closed clinic. Clinic covariates include clinic size, average physician age, share of female physicians, and share of physicians with an immigration background. Patient covariates include average patient age, share of children (age <18), share of female patients, and share of patients with an immigration background. Standard errors in parentheses are clustered at the receiving clinic level.

F.2 Placebo outcomes

Table F.3: Estimates of effects on prescribing intensity of statin

	(1) OLS	(2) IV
IV: Mean prescribing of closed clinics	0.010	
	(0.014)	
Number of prescribed patients (antibiotics)		0.008
		(0.010)
Clinic covariates	Yes	Yes
Patient covariates	Yes	Yes
Clinic FE	Yes	Yes
Year-quarter FE	Yes	Yes
Observations	31,924	31,924
Outcome mean	-2.28	-2.28
First-stage F-stat		538.45

Notes: Number of prescribed patients and the outcome, statin prescriptions per consultation, are in log-scale. Clinic covariates include clinic size, average physician age, share of female physicians, and share of physicians with an immigration background. Patient covariates include average patient age, share of children (age <18), share of female patients, and share of patients with an immigration background. Standard errors in parentheses are clustered at the receiving clinic level.

Table F.4: Estimates of effects on prescribing intensity of antidiabetic drugs (A10)

	(1) OLS	(2) IV
IV: Mean prescribing of closed clinics	-0.008	
	(0.035)	
Number of prescribed patients (antibiotics)		-0.006
		(0.026)
Clinic covariates	Yes	Yes
Patient covariates	Yes	Yes
Clinic FE	Yes	Yes
Year-quarter FE	Yes	Yes
Observations	32,197	32,197
Outcome mean	-2.28	-2.28
First-stage F-stat		543.18

Notes: Number of prescribed patients and the outcome, antidiabetic (A10) drug prescriptions per consultation, are in log-scale. Clinic covariates include clinic size, average physician age, share of female physicians, and share of physicians with an immigration background. Patient covariates include average patient age, share of children (age <18), share of female patients, and share of patients with an immigration background. Standard errors in parentheses are clustered at the receiving clinic level.